1..

| US ENVIRONMENTAL PRO ON AGENCY OFFICE OF PESTICIDES PROGRAMS | SEP 23 1986 |
|--|---|
| REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460 | TERM OF ISSUANCE |
| NOTICE OF PESTICIDE: REGISTRATION | NAME OF PESTICIDE PRODUCT |
| (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended) | Speer Stable Spray |
| NAME AND ADDRESS OF REGISTRANT (Include ZIP code) | |
| Г | ٦ |
| $\Sigma_{k}(\epsilon, a) = 0$ (4.2) | |
| Mario Artis Salit | |
| L | |
| | |
| NOTE: Changes in labeling formula differing in substance fro submitted to and accepted by the Registration Division prior product always refer to the above U.S. EPA registration numb | to use of the label in commerce. In any correspondence on this |
| On the basis of information furnished by the registrant, the at the Federal Insecticide, Fungicide, and Rodenticide Act. | pove named pesticide is hereby Registered/Reregistered under |
| A copy of the labeling accepted in connection with this Regi | stration/Reregistration is returned herewith. |
| health and the environment, the Administrator, on his motion, icide in accordance with the Act. The acceptance of any nam | or approval of this product by this Agency. In order to protect may at any time suspend or cancel the registration of a pest- e in connection with the registration of a product under this exclusive use of the name or to its use if it has been covered |
| ian ja dud – dendrij mel y rij Proth a 3(t) r)(A) an vide – hav y ar | G. A. acc. Janes A. P. C. 27 |
| 1. Submit and/or dite all data r it y us produce undo. FIFRA s d tom 30 registrant of simular produces rosub | · · · · · · · · · · · · · · · · · · · |
| z. Mak the labeling change to to the mart. | that is the many your end as the contraction |
| the off of participations | mail roll , in Morrison. |
| ha Pencambo the Tably em- proper nta | - da cambo fini sarrivo di |
| | of limius. On the Confidentia) limit: E. Prop. Hent A46 arc 22 and x they are 20 and 18. |
| = " | to read "Scourtly whap riginal of a newscapers and discard in |
| | • |
| ATTACHMENT IS APPLICABLE | |
| SIGNATURE OF APPROVING OFFICIAL | DATE |

- e. Within a year from the date on this letter submit storage stability data.
- 3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

rturo E. Castillo

Acting Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (T8-767C)

Enclosures

SPEER STABLE

Kills and repels horseflies, horn flies, stable flies, deer flies, houseflies, mosquitos, gnats, and wasps.

ACTIVE INGREDIENTS:

| Pyrethrins0.20% | |
|--|--|
| *Piperonyl Butoxide, Technical0.40% | |
| +N-Octyl Bicycloheptene Dicarboximide0.67% | |
| **Dipropyl Isocinchomeronate | |
| ***2,3;4,5-Bis(2-Butylene)Tetrahydro- | |
| 2-Furaldehyde | |
| | |

INERT INGREDIENTS:....

*Equivalent (Butylcarbityl)(6to 0.32% Propylpiperonyl) Ether and .080% related compounds.

+MGK 264 Insecticide Synergist, U.S. Pat.No. 2,576,512.

MGK Repellent 326, U.S.Pat. No. 2,757,120. *MGK Repellent 11.

For use in barns for protection of dairy cattle, beef cattle, and horses.

KEEP OUT OF REACH OF CHILDREN

CAUTION-

SEE ADDITIONAL PRECAUTIONS ON SIDE PANEL

NET WEIGHT___OZ.

MANUFACTURED BY: SPEER PRODUCTS, INC. MEMPHIS, TN 38118

EPA REG.NO.11715

EPA EST. NO. 11715-TN-1

ACCEPTED with CONTRINTS in EPA Letter Dareit

SEP23 1986

Print of the Persons becomminded bergwicks and Redemonth Aut the Remoderate few the personalise THE REST WHEN DEA THE YOU

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid inhalation of spray mist. Avoid contact with skir, eyes, or clothing. In case of contact, immediately flush contaminated skin with plenty of water. Avoid contamination of feed and foodstuffs. Do not spray directly over food or milking utensils. Wash hands with soap and water after use.

ENVIRONMENTAL HAZARDS

Remove pets, birds and cover fish aquaria before spraying. This product is toxic to fish. Keep out of lakes, streams, or ponds, Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

EXTREMELY FLAMMABLE. Contents under pressure. Keep away from fires, sparks, and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130 F may cause bursting.

STATEMENT OF PRACTICAL TREATMENT

If swallowed - Call a physician immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia.

If inhaled - Remove victim - fresh air. Apply respiration if indicated.

If on skin - Wash contaminated skin with soap and water.

If on eyes - Flush eyes with plenty of water for at leasts 15 minutes. Get medical attention immediately.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

FOR FACE FLIES: Remove mucous secretions and saliva deposits from face of animal. Hold dispenser 18 to 24 inches from animal. Spray face and other body regions for 3-4 seconds per animal. Repeat daily, as needed.

FOR HORSEFLIES, HORN FLIES, STABLE FLIES, DEER FLIES, HOUSEFILES, MOSQUITOS, AND GNATS: Hold container 18 to 24 inches from animal. Apply lightly to coat with a sweeping motion. Apply to back, belly, ear, or leg regions. Do not spray entire animal. Spray young animals sparingly. Do not apply to backs of horses prior to saddling. Mist directly over surface wounds, minor cuts, and skin abrasions to keep flies away.

FOR USE IN BARNS: Close all doors and windows and direct spray upward into center of area with a slow sweeping motion. Spray 5 to 10 seconds for each 1000 cubic feet of space. Do not remain in treated areas. Keep barn closed for 10 minutes after spraying. Ventillate thoroughly after treatment and before re-entry.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in cool place away from heat or open flame.

DISPOSAL: Replace cap and discard container

in trash. Do not incinerate or

puncture.