

Page 13

11715-182

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

JUL 2 1992

301 / 415670  
17 / 4

Mr. T. David Williams  
Speer Products, Inc.  
4242 B.F. Goodrich Blvd.  
Memphis, TN 38118

Dear Mr. Williams,

Subject: Resubmission - Revised Labeling  
Speer Multi-Purpose Insecticide Spray  
EPA Registration Number 11715-182  
Your Submission Dated May 21, 1992

The labeling referred to above, submitted September 30, 1991 in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

1. Under the instructions to treat garments (i.e., Clothes Moths) add:

Allow all treated articles to dry thoroughly before use.

2. Complete the paragraph under "ENVIRONMENTAL HAZARDS" as it appeared on the last accepted labeling as follows:

This pesticide is highly toxic to fish. Do not apply to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

Submit five copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely,

George T. LaRocca  
Product Manager (13)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

Enclosure			CONCURRENCES				
SYMBOL							
SURNAME							
DATE							

# SPEER MULTI-PURPOSE INSECTICIDE SPRAY

- \*Contains the synthetic pyrethroid PERMANONE
  - \*Controls pests up to four weeks. Controls residential, industrial and garden insects.
  - \*For use on dogs and on premises to control fleas and ticks. Aqueous, stable.
- PERMANONE- Registered Trademark of Fairfield American Corporation

## ACTIVE INGREDIENT:

*Permethrin.....	0.50%
INERT INGREDIENTS.....	99.50%
TOTAL.....	100.00%
*(3-phenoxyphenyl)methyl (+/-)cis/trans 3-(2,2-dichloroethenyl)2,2-dimethylcyclopropanecarboxylate cis/trans ratio: Min 35% (+/-)cis and max. 65% (+/-)trans	

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**  
See panel for additional precautions

Net Wt. Oz.  
Manufactured by:  
**SPEER PRODUCTS, INC.**  
4242 B.F. Goodrich Blvd.  
Memphis, TN 38118

EPA Reg. No. 11715-182 EPA EST. No. 11715-TN-1

## STATEMENT OF PRACTICAL TREATMENT

- IF INHALED:** Remove victim to fresh air. Get medical attention
- IF IN EYES:** Flush with plenty of water. Contact a physician if irritation persists
- IF ON SKIN:** Wash affected areas immediately with soap and water. Get medical attention if irritation persists.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid inhalation of spray mist and contact with skin, eyes, or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed or feedstuffs. Remove pets and birds and cover fish aquaria before space spraying or surface applications. Vacate room after treatment and ventilate before occupying. Do not allow children or pets to contact treated areas until surfaces are dry. Do not apply in pre-schools, nurseries or any classrooms when in use. Do not apply in rooms while occupied by the elderly or infirm. This product is not for use on humans. If lice infestations should occur on humans, consult your physician or pharmacist for a product for use on humans. Animals: Do not spray directly in/or eyes, mouth or genitalia of pets. Use protective gloves or mitts. Do not treat or cause exposure to puppies less than four weeks old.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish.

## PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130 degrees F. may cause bursting. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

This product is acceptable as a pesticide for use in the inedible product areas of official establishments operating under the Federal meat, poultry, shell egg grading, and egg products inspection programs.

Do not use in food/feed areas of food handling establishments, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food/feed is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food/feed processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Nonfood areas are areas such as garbage rooms, lavatories, floor drains, to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

## SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches.

Houseflies, Mosquitoes, Gnats, Wasps, Hornets, Bees, Blow Flies and Small Flying Moths: Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Repeat treatment as necessary.

Cockroaches, Waterbugs, Palmetto Bugs, Silverfish, Firebrats, Crickets, Scorpions, Spiders, Millipedes, Centipedes, Sowbugs, and Pillbugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moths, Angoumois Grain Moths: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application as necessary.

Exposed stages of Confused Flour Beetles, Saw-Toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpet. Allow carpet to dry thoroughly before use. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Allow all treated articles to dry thoroughly before use. Repeat treatment as necessary.

To control Clothes Moths: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

For control of Fleas and Ticks: For rapid control, use 1/2 oz. per 1000 cubic feet. For best results, keep animals in treated area for 24 hours.

To control Fleas: To control fleas on sheep, hogs and horses, use 1/2 oz. per 1000 cubic feet. For best results, keep animals in treated area for 24 hours.

To control Blood Sucking Insects: Repeat every 3 weeks if necessary. For Swine: Do not ship or treat.

To control Bedbugs: Repeat every 3 weeks if necessary. For control of Bedbugs, use 1/2 oz. per 1000 cubic feet.

Fleas and Ticks (Adults): Repeat every 3 weeks if necessary. For control of Fleas, use 1/2 oz. per 1000 cubic feet.

To control Fleas, Ticks and Mosquitoes: Repeat every 3 weeks if necessary. For control of Fleas, use 1/2 oz. per 1000 cubic feet.

For the control of Garden Insects: Repeat every 3 weeks if necessary. For control of Garden Insects, use 1/2 oz. per 1000 cubic feet.

May be used on plants, trees and shrubs: Repeat every 3 weeks if necessary. For control of Garden Insects, use 1/2 oz. per 1000 cubic feet.

Effective when used as a spot treatment: Repeat every 3 weeks if necessary. For control of Garden Insects, use 1/2 oz. per 1000 cubic feet.

When spraying plants, do not spray directly on foliage: Repeat every 3 weeks if necessary. For control of Garden Insects, use 1/2 oz. per 1000 cubic feet.

Directly whenever possible: Repeat every 3 weeks if necessary. For control of Garden Insects, use 1/2 oz. per 1000 cubic feet.

## STORAGE

Do not contaminate water. Storage: Store in a cool, dry place. Disposal: Wrap container in plastic and dispose of in trash. Do not reuse.

**BEST AVAILABLE**

ACCEPTED  
with COMMENTS  
in EPA Letter Dated  
JUL 2 1973