

# SPEER POINT FIVE PYRETHRIN SPRAY

## Contains Pyrenone®

For use in schools, hospitals, institutions, supermarkets, dairies, hotels, restaurants, drive-ins, motels, drug stores, food plants, milk rooms and animal quarters.

Provides rapid control of houseflies, mosquitoes, small flying moths (Millers), gnats, wasps and accessible stages of cockroaches, water bugs, silverfish, flour beetles, weevils, grain mites and spiders.

Kills and repels stable flies, horse flies, face flies, deer flies, house flies, mosquitoes and gnats.

Controls garden insects on ornamental and vegetable plants.

## ACCEPTABLE FOR USE TO DAY OF HARVEST ON EDIBLE PLANTS

### ACTIVE INGREDIENTS:

Pyrethrins ..... 0.5%

Piperonyl butoxide, technical\* ..... 5.0%

INERT INGREDIENTS: ..... 94.5%  
100.0%

\* Equivalent to 4% (butylcarbityl)(6-propyl-piperonyl) ether and 1% related compounds.

Pyrenone® - Registered trademark of Roussel Uclaf Corporation.


## KEEP OUT OF REACH OF CHILDREN

## CAUTION

See side (back) panel for additional precautions.

NET WEIGHT: \_\_\_\_\_

EPA Registration Number: 11715 - 257  
EPA Establishment Number: 11715-TN-1



## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Do not apply directly to food. In commercial food handling facilities, do not apply while food processing is underway. Cover or remove any food or food processing surfaces and equipment prior to application. After space spraying in food processing areas, wash all equipment, benches, shelving, etc. where exposed food will be handled with an effective cleaning compound followed by a portable water rinse prior to use. All food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

Remove pets, birds, and cover fish aquariums before spraying.

### FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-757-4943 for emergency medical treatment information.	

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

### PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F. may cause bursting.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### SHAKE BEFORE USING AND AT INTERVALS DURING PROLONGED USE.

House flies, mosquitoes, gnats, wasps and small

11715-257

6/19/2003.

Page 182

**flying moths (millers):** Close all doors and windows and spray room at the rate of 2 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat as necessary.

**Cockroaches, waterbugs, silverfish, crickets, spiders, centipedes and sow bugs:** Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floor, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

**Ants:** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

**Fleas:** Spray sleeping quarters, bedding, floor and floor coverings where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

**Carpet beetles:** Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

**Bedbugs:** Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary.

**In animal quarters and milk rooms:** For rapid control of house flies, mosquitoes, gnats, wasps and small flying moths, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

**To treat animals:** To protect beef cattle, dairy cattle and horses from attacks of stable flies, horse flies, deer flies, house flies, horn flies, mosquitoes and gnats, spray about 3 seconds on each side being careful to spray back, withers and forelegs

thoroughly. To protect from face flies: spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

**Exposed stages of indian meal moths and angoumois grain moths:** Close doors and windows. Direct spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for that purpose.

**Exposed stages of flour beetles, weevils and grain mites:** Spray into cracks, and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary. Infested stored products in the home should be disposed of.

**Controls garden insects on ornamental vegetable plants:** Spray with a slow sweeping motion, not closer than 18 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants spray in short bursts. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90° F. Contact insects directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rains.

**Acceptable for use to day of harvest on edible plants.**

**Use on ornamentals (roses, asters, dahlias, african violets, azaleas, euonymus, hydrangeas, junipers, lilacs)** to control aphids, rose chafers, japanese beetles, rose slugs, tent caterpillars,

scales (including euonymus and juniper scales), leafhoppers, tarnish plant bugs, lace bugs (including azalea lace bugs), thrips, (exposed), mealy bugs (including citrus mealy bugs), mites (including red spiders, spiders and broad mites), plant bugs, white flies, leaf miners, hydrangea leaf tiers, bagworms and leaf rollers.

**Use on vegetables (peppers, eggplants, potatoes, cucumbers, squash, melons, beans sweet corn and tomatoes)** to control flea beetles, hornworms, stink bugs, colorado potato beetles, aphids (including green peach aphids), leafhoppers, blister beetles, striped and spotted cucumber beetles, mexican bean beetles, white flies, 12-spotted beetles and japanese beetles.

#### STORAGE AND DISPOSAL

**STORAGE:** Store in a cool, dry area away from heat or open flame. Do not transport or store below 32°F.

**DISPOSAL:** Do Not Puncture or Incinerate!

**If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

[Distributors may choose their own emergency number and time of operation.]

**MANUFACTURED BY:  
SPEER PRODUCTS, INC.  
4242 B. F. GOODRICH BLVD.  
MEMPHIS, TN 38118**

c:\mispub\reg257.pubNOTIFICATION[08/13/01]-11715-257-  
NOTIF-2001-1&6(4-17-03)

272