SPEER FIRE ANT BAIT

[] or () indicates alternate or optional wording

[Speer Fire Ant Bait] is an insect growth regulator similar to the naturally occurring insect growth hormones which interferes with [stops] their growth or development

[Speer Fire Ant Bait] inhibits worker replacement in fire ant colonies

[Speer Fire Ant Bait] kills [controls] developing immature fire ants and degenerates the reproductive organs of the queen

[Speer Fire Ant Bait] provides a shift in caste differentiation from worker fire ants to sexual forms

[Speer Fire Ant Bait] reduces the quantity of the worker brood within two weeks

[Speer Fire Ant Bait] reduces the colony size by 85% in one month and by 96% in two months

[Speer Fire Ant Bait] causes total colony death within six months

Effective in the control of the Red Imported Fire Ant [Solenopsis invicta] [and other Solenopsis species] and the Big-Headed Ant

Easy to use granular formulation

Fire ants seek out the [Speer Fire Ant Bait] and take it back to the colony

[Speer Fire Ant Bait] breaks the reproductive life cycle of the fire ant causing starvation of the colony

Worker fire ants bring the bait into the mound, feed it to the queens, reducing egg production leading to colony death

ACTIVE INGREDIENT

KEEP OUT OF REACH OF CHILDREN CAUTION

NET CONTENTS:

EPA REG. NO. 11715-311 EPA EST. NO 11715-TN-1

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid breathing vapors or spray mist. Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact, flush with plenty of water. Wash with soap and warm water after use. Obtain medical attention if irritation persists. Avoid contamination of food or feedstuffs.

DIROT AIR	
	FIRST AID
IF SWALLOWED:	Call poison control center or
SWALLOWED:	doctor immediately for treatment advice.
	Have person sip a glass of
	water if able to swallow.
	Do not induce vomiting
	unless told to do so by the
	poison control center or
	doctor.
	Do not give anything by
	mouth to an unconscious
	person
IF INHALED:	Move person to fresh air.
	If person is not breathing, call
	911 or an ambulance, then
	give artificial respiration,
	preferably by mouth-to-
	mouth, if possible.
	Call a poison control center or
	doctor for further treatment
IF ON SKIN OR	advice.
CLOTHING:	Take off contaminated clothing.
CLOTHING:	Rinse skin immediately with
	plenty of water for 15-20
	minutes.
	Call a poison control center or
	doctor for treatment advice.
	ACCEPTED
	WOOMLIND
	JUL 2 1 2003
	Under the Federal Insecticide,
	Fungicide, and Rodenticide Act,
!	as amended, for the pesticide registered under
	RPA Reg. No. /17/5-3/1

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or quotor for treatment advice.

HOT LINE NUMBER

Have the product container (r label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-757-4943 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

[Speer Fire Ant Bait] is a slow acting, highly effective fire ant control product. Fire ant workers pick up the bait granules and take it back to the colony. Within 3 to 4 weeks there is substantial colony mortality and within eight weeks the majority of the colony population has been eliminated.

Worker fire ants bring the [Speer Fire Ant Bait] into the mound and feed it to both the queen and the immature fire ants, thereby preventing worker replacement. Due to degeneration of reproductive organs, the fire ant queen cannot replace workers. The lack of worker replacement results in colony death as the existing worker ants age and die.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ALL DIRECTIONS COMPLETELY BEFORE USE.

For best results, follow directions for specific use areas.

[Speer Fire Ant Bait] is intended to be applied by lawncare operators and landscape personnel for outdoor

use to control red imported fire ants in the following sites.

Agricultural sites defined as sod farms and commercial

nurseries producing turf and ornamentals for resale

for seed.

Uncultivated, nonagricultural areas including airports,

roadsides, highway rights-of-way, utility rights-of-way,

campgrounds and industrial sites.

Institutional sites defined as areas around properties

or

facilities providing a service to public or private organizations including, but not limited to, hospitals, nursing homes, schools, universities, museums, libraries.

sport facilities (golf courses), and office buildings.

Residential/domestic sites, defined as areas associated

with the household or home life including, but not limited to, apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions.

hospitals or convalescent homes.

WHEN TO APPLY:

[Speer Fire Ant Bait] should be applied in the early Spring or Summer at the first sign of fire ant activity. Application may be made any time of the day, but is more effective when ants are actively foraging, usually when the soil temperature is above 60°F. Avoid application if rain is expected within 4 to 6 hours. Do not water treated area for 24 hours after application. In the event of prolonged rains after initial treatment, reapplication may be necessary.

In areas where large mounds remain active, areas of heavy infestation or reinfestation occurs, a second application of [Speer Fire Ant Bait] may be advisable after 12 to 16 weeks to maintain sufficient levels of active ingredient. In very large mounds, some worker fire ants may persist for a while after the queen is dead. For 7-10 days after treatment, do not apply any other fire ant pesticide. This is to allow sufficient

time for the worker ants to seek out the bait and take it

back to the mound. After 14 days, it may be desirable to apply a registered fire ant pesticide to kill remaining foraging ants.

INDIVIDUAL MOUND TREATMENT:

Do not disturb the mound. Do not apply to top of mounds. Apply 1 to 4 level tablespoons of [Speer Fire Ant Bait] per mound, uniformly distributing material 3 to 4 feet around the mound. Retreat after 12 to 16 weeks if needed.

BROADCAST OR AERIAL APPLICATION:

Apply uniformly with equipment calibrated to give the correct dosage. Apply at 1 to 1.5 lb. of product per acre or the equivalent rate of 0.35 to 0.5 oz. per 1000 sq. ft.. Retreat after 12 to 16 weeks if needed.

BIG-HEADED ANT (Pheidole megacephala) CONTROL

Apply uniformly with ground or hand-held equipment calibrated to deliver the correct dosage. Apply at 1 to 1.5 lb. of product per acre (or 0.35 to 0.5 oz. per 1000 sq. ft.). Treat surrounding ares, adjacent borders, and fence lines to help prevent ant migration into the treated areas. Repeat after 12 to 16 weeks if needed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool, dry place. Keep container closed.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved State and Local procedures.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

MANUFACTURED BY: SPEER PRODUCTS, INC. 4242 B. F. GOODRICH BLVD. MEMPHIS, TN 38118

C:\MSPUB\REG311(07/14/98)11715-311-NOTIF-2001-1(5-1-03)