

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460 EPA Reg. Number:

Date of Issuance:

11715-317

APR 23 1997

NOTICE OF PESTICIDE:

x Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance: Conditional

Name of Pesticide Product:

NeoPerm Industrial Insect Killer

Name and Address of Registrant (include ZIP Code):

Speer Products, Inc.

P.O. Box 18993

Memphis, TN 38181-0993

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number:

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA sec. 3(c)(5) or FIFRA sec. 4 when the Agency requires all registrants of similar products to submit such data.
- 2. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 11715-317".
- 3. To comply with PR Notice 96-6 issued October 1, 1996, make the following changes to the label:
 - a. At the top of the rear or side panel insert the phrase "READ ENTIRE LABEL BEFORE EACH USE.".
 - b. Change the statement "To control fleas, ticks, and lice on dogs" to "To control fleas, ticks, and lice on dogs ONLY".

Date

4/23/87

SIM

George T. LaRocca, Product Manager (13)
Insecticide-Rodenticide Branch
Registration Division (7505C)

Page Two 11715-317

c. Delete the Statement "Not recommended for use on puppies less than four weeks old". Replace with the following:

"Do not use on puppies less than 12 weeks old. Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant or nursing animals.".

en da un antario de la marcina de la casación de la región de conservación de la conservación de la companya d

THE REPORT OF THE PROPERTY OF

d. Under the precautionary statements add the following:

"Sensitivities may occur after using any pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately."

4. Submit two (2) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec.

6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

SPEER NEOPERM INDUSTRIAL INSECT KILLER

Contains Pyrenone* and Permanone* Controls Industrial Insects Kills on Contact For Commercial/Agricultural Use A Broad Spectrum Insecticide Highly Effective Against Ants (Including Carpenter Ants), Bees, Beetles (Including Carnet, Flour, Grain, and Ground Beetles), Centipedes, Clover Mites, Cockroaches, Crickets (Including Mole Crickets), Earwigs, Firebrats, Fleas, Millinedes, Moths, Pillbugs, Scorpions, Silverfish, Sowbugs, Spiders, Ticks, Wasps, and Other Pests Listed on the Label, Kills Lice and Their Eggs For Use on Bedding, Furniture and Other Inanimate Objects Infested with Lice Non-Staining No Unpleasant Odor Water-based Formula For Use on Dogs and on Premises to Control Fleas and Ticks. Controls Fleas and Ticks on Dogs for 14 Days.

ACTIVE INGREDIENTS:

710-11-14	(11 1 U 1 1 L	4.4.2 1.1.3 L L	Q,			
Pyrethrin:	5. , ,		, ,			0.10%
Piperony!	Butoxid	e, Techi	nical*.	.,,,,,		0,50%
						0,20%
INERT I	NGRED	IENTS	:. , ,			99,20%
						other and 0.1%
related comp	pounds.			:		4) 1
						nethonyl) 2.2
dimethyleye	lopropanec	arboxy late	. Cistre	nis tatio:	min. 35% (+	/-) cls and max
65° \ (-/-) #	ans.	1 1		.		
Рутепопе ап	d Permano	ne Regis	tered Tra	ademarka	of AgrEvo I	in ironmental
Health.	1 1		7 1 1		- 44	**
, ,			100		·	

EPA REG. NO. 11715-not yet assigned G7 R EPA EST. NO. 11715-TN-1

KEEP OUT OF REACH OF CHILDREN ACCEPTED CAUTION with COMMENTS in EPA Letter Bete Side Panel For Additional Precautions

Under the Federal Insecticide, Fungicide, and Rodenticide Act as smended, for the pesticide registered under EPA Reg. No.

STATEMENT OF PRACTICAL TREATMENT

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if inhaled, Avoid breathing vapor or spray mist, Remove contaminated clothing and wash before reuse. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before spraying. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of Insects. Repeat treatment as necessary. equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open temperatures above 130°F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner where possible. Repeat as necessary. inconsistent with its labeling.

rooms, but excluding areas where foods may be prepared or necessary. held.

floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Not for use in USDA meat and poultry plants.

Do not use in rooms/areas while occupied by patients, the elderly or infirm. Do not apply to classrooms when in use.

SHAKE WELL JUST BEFORE USING.

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches.

INDOORS:

Cluster Flies, Houseflies, Fruit Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees, Black Flies, Skipper Flies, Blow Flies and Small Flying Moths: Close all doors and windows and direct spray upward into center of room with a slow sweeping motion. Spray 5-10 seconds for average room, moving toward the door. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Leave room and keep area closed for 15 minutes after spraying. Ventilate room thoroughly before re-entry. Sweep up and discard fallen

Cockroaches, Waterbugs, Palmetto Bugs, Silverlish, Firebrats, Crickets, Scorpions, Spiders, Millipedes, Centipedes, Sowbugs and Pillbugs: Spray into hiding places flame. Do not puncture or incinerate container. Exposure to such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants:

Indian Meal Moths, Angoumois Grain Moths: Close doors Do not use in food/feed areas of food/feed handling and windows direct spray into all parts of room especially establishments, restaurants or other areas where food/feed is around stored food product containers, pallets and darkened commercially prepared or processed. Do not use in serving areas. Spray 7 to 10 seconds for each 1000 cubic feet of space, areas while food is exposed or facility is in operation. Serving moving toward the door. Leave room and keep area closed for areas are areas where prepared foods are served such as dining 15 minutes following treatment. Repeat application as

Non-food areas are areas such as garbage rooms, lavatories, Exposed stages of Confused Flour Beetles, Cigarette Beetles, Drug Store Beetles, Saw-Toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites in exposed adult and larvae stages: Spray into cracks and crevices, processing and handling equipment, around cartons of containers or other areas where these insects tend to congregate. Contact as many insects as possible. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Space spray: Spray

Leave room and keen area closed for 15 minutes following four weeks old. treatment. Repeat application weekly or as necessary. Infested stored food products should be disposed of, or treated using an TO KILL Lice and Louse eggs: Spray in an inconspicuous Point spray nozzle away from face. Hold can on slight approved product intended for this purpose.

Carpet Beetles: Spray directly on beetles, floors, baseboards. Hold container upright with nozzle away from you. Depress wind if breeze is blowing. shelves and carpets. Repeat treatment as necessary.

seams. Take beds apart and spray into all joints. Treat be either laundered or dry cleaned. Allow all sprayed articles drift over area. Direct spray toward bushes, or grass from at baseboards; moldings and floors. Repeat treatment as to dry thoroughly before use. necessary. Allow all sprayed articles to dry thoroughly before

TO CONTROL Clothes Moths and Carpet Beetles on wool Bees and Small Flying Moths, close windows and doors and insects and their nests from approximately 2 feet away. and other proteinaceous fibers. Remove infested articles from spray at the rate of 2 seconds per 1000 cubic feet of space. For Contacted insects will fly away from fog. Application should storage, brush thoroughly, and air for several hours in sunlight, best results, keep area closed for 10 minutes after spraying. Do be made in late evening when insects are at rest. Spray into if possible. Apply spray liberally to empty chests, closets, not remain in treated area and ventilate after treatment is hiding and breeding places contacting as many insects as bureaus and other storages directing the spray into cracks, completed. joints and crevices. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet. To Treat Livestock: To protect beef cattle, dairy cattle, goats, CRAWLING INSECTS: Ants, Cockroaches, including from fabric to avoid staining.

floor and direct spray toward floor and lower walls at a rate of being careful to spray back, withers and forclegs thoroughly, and chairs to repel ants. Spray ant hills for best control. 10 seconds per 100 sq. ft, making sure that all floor, sofa and To protect from Face Flies, spray the face and head, but do not chair surfaces are contacted. Leave room and keep area closed spray into eyes. Repeat treatment when flies are troublesome. for 15 minutes. Open and ventilate before re-occupying, Repeat treatment after 7 days or as needed.

Fleas & Ticks (Adults and Larvae): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, TO CONTROL Blood Sucking Lice: Apply to the infested entire floor and floor covering, with special attention to cracks areas of the animal, using a stiff brush to get the spray to the Buyer assumes all risks of use, storage or handling of this and crevices around baseboards, and window and door frames, base of the hair. Repeat every 3 weeks if required. Infested upholstered furniture and mattresses may also be treated with this product. A test application should be made to TO CONTROL Poultry Lice: Spray roosts, walls and nests or upholstery or drapery fabrics in an inconspicuous place before cages thoroughly. This should be followed by spraying over use.

TO CONTROL Fleas, Ticks and Lice on Dogs: Starting at FOR CONTROL OF Bedbugs and Mites in poultry houses: the head, stroke against the lay of the hair, spraying the parted Spray crevices of roost poles, cracks in walls, and cracks in fur in short bursts, directly behind the hand to insure nests where the bedbugs and mites hide. penetrating the coat. Dampen the fur back to the tail. Repeat application weekly if necessary to control infestations.

20 seconds per 1000 cu. ft. of space, moving toward the door. NOTE: Not recommended for use on puppies less than

area to test for possible staining or discoloration. Inspect again downward angle. For best results, spray when air is calm, after drying, then proceed to spray entire area to be treated. Allow a few minutes for product to take effect. Spray with valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray only those garments and FLYING INSECTS: Flies, Mosquitoes, Gnats, Small Flying Bedbugs: Spray mattresses, particularly around tufts and parts of bedding, including mattresses and furniture that cannot Moths, Hornets, Wasps and Yellow Jackets: Allow fog to

> IN ANIMAL QUARTERS AND MILK ROOMS: For rapid control of House Flies, Mosquitoes, Gnats, Wasps, Hornets, Mud Daubers and Wasp Nests: Spray mist directly on

Fleas in Buildings - Space spray: Hold can 36 inches above Mosquitoes and Gnats: Spray about 3 seconds on each side hitting as many insects as possible. Also spray legs of tables

last treatment.

the birds with a fine mist.

OUTDOORS:

least 4 to 6 feet away. Give special attention to breeding areas such as garbage cans and compost piles. Repeat as necessary,

possible.

sheep, hogs and horses from attacks of Stable Flies, Horse Asian Cockroaches, Crickets, Mole Crickets, Centipedes Flies, Deer Flies, Face Flies, House Flies, Horn Flies, and Silverfish; Spray infested surface of patio or picnic area,

STORAGE AND DISPOSAL

Store in cool dry area away from children. Do not transport or For Swine: Do not ship animals for slaughter within 5 days of store below 32°F. Wrap container in several layers of newspaper and dispose of in trash. Do not inclnerate or puncture.

material not in strict accordance with directions given herewith.

MANUFACTURED BY SPEER PRODUCTS MEMPHIS, TN 38118

AUG 13 P1:42

KECD EPA/OPP/DP01