SPEER PERMETHRIN **DUSTING POWDER**

Contains Permanone

Provides Ouick Killing Action and Residual Control of Fleas and Ticks on Dogs for up to 14 Days

A Broad Spectrum Ready-To-Use Insecticide Highly Effective Against Ants, Carpenter Ants, Bees, Carnet Beetles, Centipedes, Cockronches, Crickets, Eurwigs, Firebrats, Fleas, Flour Beetles, Grain Beetles, Ground Beetles, Millipedes, Mole Crickets, Moths, Pillbugs, Scorpions, Silverfish, Sowbugs, Spiders, Ticks, Wasps, and Other Pests as Listed on Side Panel. Odorless, Nonstaining Formulation.

For Household Use Only: For Use in Rooms, Apartments, Homes, Attics, Basements, Campers, Boats, Household Storage Areas, Garages, Pet Sleeping Areas, Cabins,

ACTIVE INGREDIENTS:

Piperonyl Butoxide, Technical*	2.00%
Permethrin**	
INERT INGREDIENTS:	<u>97.50%</u>
	100.00%


- *Equivalent to 1.6% (butylearbityl)(6-propylpiperenyl) ether and 0.4% related
- **(3-phenoxyphenyl) methyl (±) cis/trans 3-(2,2-dichloroethenyl) 2,2dimethylcyclopropanecarboxylate. Cis/trans ratio; min. 35% (±) cis and max. 65% (±) trans.

PERMANONE - Registered Trademark of Aprilyo Environmental Health, Inc.

KEEP OUT OF REACH OF CHILDREN CAUTION

See Side Panel For Additional Precautionary Statements

NET	' ^^	NT	RN1	ΓÇ



READ ENTIRE LABEL BEFORE EACH USE ON ANIMAL APPLICATION FOR DOGS ONLY

PRECAUTIONARY STATEMENTS **ILAZARD TO HUMANS & DOMESTIC ANIMALS** CAUTION

Harmful if inhaled, Avoid breathing dust. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Take care not to introduce the product into the air during indoor application; use only in well ventilated areas. Do not apply where children are likely to contact treated surfaces. Remove or cover aquaria before application. Do not apply to areas, surfaces or utensils which will contact food or feed. Exposed food should be covered or removed during application. All food processing surfaces and utensils should be covered during application or thoroughly washed with soap and water after use.

Do not apply product directly in/on eyes, mouth or genitalia of dogs. Do not treat or cause exposure to puppies less than twelve weeks old. DO NOT use this product on cats. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals, or animals on medication. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

FIRST AID

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-tomouth. Get medical attention.

ENVIRONMENTAL HAZARDS

disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a found. manner inconsistent with its labeling.

INDOOR USE:

applications with hand or power duster (Actisol® or similar deemed necessary to avoid stings. Using hand or power

equipment), or by other suitable means, to hiding and runway areas and other places where insects are found. To apply insecticide directly into cracks and crevices, use a bulbous duster or other suitable equipment. Apply lightly and uniformly to infested areas.

COCKROACHES (including adult and immature stages of both non-resistant and organophosphate and carbamate resistant strains), ANTS, CRICKETS, SILVERFISH AND FIREBRATS: Apply thoroughly at a rate of 3 to 6 ounces per 100 square feet to all areas where these pests crawl and hide. especially in cracks and crevices and hidden surfaces around sinks and storage areas, behind baseboards, around doors and windows, behind and under refrigerators, cabinets, sinks and stoves and in attics and crawl spaces. For Auts, sprinkle freely around ant trails, doors and windows and other places where ants enter premises around nests hidden under steps. brickwork, concrete, etc. Break apart accessible nests and scatter powder freely on and around debris. Repeat as necessary.

BROWN DOG TICKS AND FLEAS IN BUILDINGS:

Apply dust at a rate of 3 to 6 ounces per 100 square feet to infested areas such as pet beds and resting quarters, nearby cracks and crevices, under the edges of rugs and floor coverings, between and under cushions of upholstered furniture and other areas where these pests may be present. Old bedding of pets should be removed and replaced with clean fresh bedding after treatment of pet areas.

SPIDERS: Apply at a rate of 3 to 6 ounces per 100 square feet along and behind baseboards, to window and door frames, corners, pipes, storage localities, attics, crawl spaces and other areas over which these pests may crawl.

SOWBUGS, PILLBUGS, GROUND BEETLES. This product is toxic to fish and other aquatic organisms. Do EARWIGS, MILLIPEDES, CENTIPEDES AND not apply directly to water. Do not contaminate water when SCORPIONS: Apply at a rate of 3 to 6 ounces per 100 square feet around doors and windows and other places where these pests may enter premises. Treat baseboards, storage areas and other locations where these pests are

WASPS AND BEES: It is generally advisable to treat wasp and bee nests in the evening when the insects are less active Speer Permethrin Dusting Powder is intended for and have returned to the nest. Wear protective clothing if

necessary, thoroughly dust nest and entrance and surrounding powder. Take care not to get the powder in animal's eyes, reused, dispose of in same manner, areas where insects light. Nests in wall voids can be located nose, mouth or genitalia. Dust bedding, kennel, sleeping by listening with your caregainst the wall. Drill a hole in the areas, carpeting and furniture used by infested animals. Do area, blow dust in, and rescal. For ground wasps, dust not treat dogs under 12 weeks of age. Repeat treatment at CONTAINERS ONE POUND AND SMALLER: Do not entrance and surrounding areas. For best results, check nests carefully one or two days after treatment to ensure complete kill, then remove and destroy nest to prevent OUTDOOR USE emergence of newly-hatched insects. If removal is not PERIMETER TREATMENT: Use Speer Permethrin

PANTRY PESTS - GRAIN BEETLES (RUSTY MERCHANT AND SAW-TOOTHED), FLOUR BEETLES, (RED AND CONFUSED), DRUGSTORE BEETLES, RICE WEEVILS, LESSER GRAIN BORERS, SPIDER BEETLES AND TOBACCO MOTHS: Treat exposed stages. Apply dust at a rate of 3 to 6 ounces per 100 square feet in cracks and crevices behind and under cupboards and cabinets for residual control. Do not apply directly on food, utensils or food storage areas.

to treat overhead nests.

CARPET BEETLES (BLACK, FURNITURE AND VARIED): Dust at a rate of 6 ounces per 100 square feet on and under edges of floor coverings, under rugs and furniture and in closets or other localities where these insects are found.

BEDBUGS: Dust mattresses at a rate of 3 ounces per 100 square feet,, particularly around tufts and seams, Take beds apart and dust into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary.

WEBBING CLOTHES MOTHS: Use this product as a Do not contaminate food, feed or water by storage or crack and crevice treatment in closets and other storage areas disposal. where these pests are found. This is an adjunct treatment and will not control moth larvae already on the clothes. Do not Pesticide Storage and Spill Procedures: Store upright at apply to clothing.

ON DOGS ONLY: Speer Permethrin Dusting Powder provides quick killing action and long lasting control of fleas and ticks on dogs.

TO KILL FLEAS, TICKS AND LICE AND TO REDUCE ITCHING AND SCRATCHING DUE TO INSECT BITES Apply powder liberally over the entire body, beginning at the head and working back. Work dust into hair and around the feet, pads and between the toes by rubbing against the lay of

weekly intervals if necessary.

feasible, retreat the nest, if necessary. Do not use this product Dusting Powder as a residual treatment for control of Ants, Bees, Brown Dog Ticks, Centipedes, Cockroaches, Crickets, Earwigs, Firehrats, Fleas, Ground Beetles. In case of emergency or for product use information please Millipedes, Pillbugs, Scorpions, Silverfish, Sowbugs, call 901/362-1950 Monday thru Friday CST (Distributors Spiders, Ticks and Wasps, Make a residual treatment around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces and other areas where pests hide.

> To help prevent invasion of buildings and structures by the above pests, treat a band of soil and vegetation 6 to 10 feet wide around and adjacent to the building or structure at a rate of 3 to 6 ounces per 100 square feet. Also treat the building foundation and crawl spaces where pests are active and find entrance. Apply thoroughly and uniformly to the band area. Pay particular attention to ant hills and wasp nests in the band area. Wasp and bee nests in trees and shrubs may be treated as noted above.

> Notice: Buyer assumes all responsibility for safety and use not in accordance with directions.

STORAGE AND DISPOSAL

room temperature. Keep away from moisture. Sweep up and dispose of with chemical waste.

Pesticide Disposal: Pesticide that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

Container Disposal: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and

duster or other suitable means, with extension tubes if the hair until dust penetrates to skin. Contact ticks with local authorities. If drum is contaminated and cannot be

reuse container. Wrap container in several lavers of newspaper and discard in trash.

may use their own phone numbers and hours of operation).

MANUFACTURED BY: SPEER PRODUCTS, INC. P. O. Box 18993 Memphis, TN 38181-0993

E:\MSPUB\reg319.pub-(10/16/97)