
 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration</p> <p>(under FIFRA, as amended)</p>	<p>EPA Reg. Number: 11715-336</p>	<p>Date of Issuance: MAY 31 2001</p>
	<p>Term of Issuance: Conditional</p>	
	<p>Name of Pesticide Product: SPI Permethrin Home & Garden Insect Killer</p>	
<p>Name and Address of Registrant (include ZIP Code): Speer Products, Inc. 4242 B.F. Goodrich Blvd. Memphis, TN 38118</p>		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is <u>conditionally</u> registered in accordance with FIFRA sec. 3(c)(7)(A), provided you:</p> <ol style="list-style-type: none"> Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. Make the following label changes: <ol style="list-style-type: none"> Revise the EPA Registration Number to read, "EPA Reg. No. 11715-336". Revise the Data Matrix associated with this product to include product performance (efficacy) studies being used to support the public health pests on the label (i.e., fleas, ticks, lice, bedbugs, spiders, scorpions, and cockroaches). 		
<p>Signature of Approving Official:  George T. LaRocca, Product Manager 13</p>	<p>Date: MAY 31 2001</p>	


2/17

- c. Under TO CONTROL FLEAS AND TICKS, include the statement, "Remove pets prior to treatment".
 - d. The retreatment statements should include a maximum frequency of application, such as, "...but not more than __ times per __". Amend each retreatment statement on the label.
 - e. Under ON LIVESTOCK, include the statement, "Avoid contact with animal's genitalia". "Do not spray into eyes or face".
 - f. We prefer you move the precautions for dogs on the front panel to the DIRECTIONS FOR USE on dogs.
 - g. The product name indicates use in/around home and gardens, however directions for use are given for livestock use. Revise the name similar to "SPI Home, Garden and Livestock Insect Killer" or "SPI Insect Killer".
 - h. Under DIRECTIONS FOR USE add, "Not for use in commercial food/feed handling establishments, restaurants or other establishments where food/feed is commercially prepared, processed or served".
3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,



George T. LaRocca
Product Manager (13)
Insecticide Branch
Registration Division (7505C)

Enclosure

SPI PERMETHIN HOME & GARDEN INSECT KILLER

FOR DOMESTIC USE BOTH INDOORS AND OUTDOORS
FOR LIVESTOCK AND PREMISE USE
READY TO USE FOR CONTROL OF HOUSEHOLD PESTS
DILUTE FOR USE ON VEGETABLES AND FRUIT AND NUT TREES
FOR CONTROL OF FLEAS, BROWN DOG TICKS AND LICE ON PREMISES AND ON DOGS
CONTROLS FLEAS AND TICKS ON DOGS FOR 14 DAYS
FOR CONTROL OF PESTS ON LIVESTOCK
WITH COUNTRY FRESH FRAGRANCE
PROVIDES 60-DAY (TWO MONTHS) CONTROL OF COCKROACHES
PROVIDES 60-DAY (TWO MONTHS) CONTROL OF STORED PRODUCT PESTS

ACTIVE INGREDIENTS:

Permethrin.....0.25%

OTHER INGREDIENTS:.....99.75%
100.00%

*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate. Cis/trans ratio: min. 35% (1/-) cis and max. 65% (1/-) trans.

ACCEPTED KEEP OUT OF REACH OF CHILDREN
WITH COMMENTS CAUTION
In EPA Letter See Side Panel for Additional Precautions
MAY 31 2001

NET CONTENTS: _____

EPA REG. NO. 11715-
EPA EST. NO. 11715-TN-1

READ ENTIRE LABEL BEFORE EACH USE
PET USE APPLIES TO USE ONLY ON DOGS

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals CAUTION

Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and food-stuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

Do not use on dogs under twelve (12) weeks old. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or other commercial use, or for commercial seed production or for research purposes.

SHAKE WELL BEFORE USING

INDOOR APPLICATION

HOUSEHOLD PESTS: For spot application only. Do not use as a space spray indoors.

Ready to use. Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

SURFACE SPRAYING: FOR 60 DAYS (TWO MONTHS) CONTROL OF: COCKROACHES, WATERBUGS, PALMETTO BUGS, SCORPIONS, MILLIPEDES, CENTIPEDES, SOWBUGS, PILLBUGS, ANTS, SILVERFISH, FIREBRATS, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES AND CADELLES: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS (ADULTS AND LARVAE): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and be-

3/4

hind baseboards, moldings, window and door frames, and entire areas of floor and floor coverings. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping the nozzle at least three feet from fabric to avoid staining.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

LIVESTOCK PESTS

Ready to use. Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from **horn flies, face flies, house flies, mosquitoes and gnats:** Apply a light mist sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES:

Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOOD-SUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE: Spray roosts, walls and nest or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

TO CONTROL BEDBUGS AND MITES IN POULTRY HOUSES: Spray crevices of roost poles, cracks in walls, and cracks in nests where the Bedbugs and Mites hide.

ON DOGS: TO CONTROL FLEAS, TICKS AND LICE: Ready to use. Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container. Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and under body. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Repeat application every two weeks if necessary. **DO NOT USE ON CATS OR KITTENS.**

OUTDOOR APPLICATION

TO HELP CONTROL TICKS IN OUTDOOR ACTIVITY AREAS: Ready to use. Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container. Treat a band of vegetation and leaf litter 6-10 feet wide and

adjacent to activity areas, especially where dense vegetation occurs.

PESTS ON THE OUTDOOR SURFACE OF BUILDINGS: To temporarily control **House Flies, Mosquitoes, Cluster Flies, Gnats and Flying Moths.** For outdoor use only as an aid in reducing annoyance from these insects.

Dilute before using. Use clean sprayer. Carefully measure and mix 1 fluid ounce of SPI PERMETHIN HOME & GARDEN and 12 fluid ounces of water. Using dilute mixture, spray pests on outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate. Repeat as necessary. Flush sprayer with clean water after each use.

ORNAMENTAL FLOWER GARDENS

Ready to use. Affix trigger, pump or other type of sprayer to container for application. Do not transfer to another container.

For use on roses, dahlias, asters and other ornamentals to kill **Japanese Beetles** hit by spray as well as **Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites and Leafminers.** May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering jews and zinnias.

4/7

Effective when used as directed against **Whiteflies, Aphids, Spiders and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms.**

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90° F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

ORNAMENTAL TREES AROUND THE HOME:

Dilute before using. Use clean sprayer. Carefully measure and mix 1 fluid ounce of SPI PERMETHIN HOME & GARDEN and 12 fluid ounces of water. Using dilute mixture, use sufficient spray to obtain full coverage of all foliage. Avoid heavy wetting. Use to control: **Pine Moths, Pine Needle Miners, Pine Beetles and Needle Scales** on pine; **Budworms** on conifers; **Oakworms** on Oak; **Fall Webworms** on cherry, elm and honeysuckle; **Aphids** on tulip poplar; **Citrus black flies** on non-bearing citrus; **Tussock Moths** on Douglas and grand firs; **Leafminers** on birch; **Elm Leaf Beetles** and **Elm Spanworms** on elms. Flush sprayer with clean water after each use.

FOR HOME VEGETABLE GARDENS: Dilute before using. Use clean sprayer. Carefully measure and mix 1 fluid ounce of SPI PERMETHIN HOME & GARDEN and 12 fluid ounces of water. Spray upper and lower leaf surface as directed. Use sufficient spray to obtain full coverage of all foliage, but do not spray to the point of runoff. Apply no more than 13 ounces of dilute mixture over 100 linear feet of row (13 ounces of diluted mixture per 100 square feet.) Flush sprayer with clean water after each use.

VEGETABLE	TARGET PESTS	REMARKS
Asparagus (pre-harvest)	Cutworms Asparagus beetle	Spray plants as needed to ensure uniform coverage. Do not make more than 4 applications. Can be applied up to 3 days prior to harvest.
Asparagus (post-harvest)	Larval stages of asparagus beetle; tarnished plant bugs, lygus bugs and adult Japanese beetles.	Treat fern or bush growth after spear harvest when insects are present.
Celery- see leafy vegetables		
Cole Crops: Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Beet armyworm; diamond back moth; imported cabbageworm, cabbage aphid, cabbage looper	Apply every 5 days or as needed. Can be applied up to one day prior to harvest. Do not make more than 10 applications on cabbage and 8 applications on the other cole crops.
Corn- see sweet corn		

Cucurbits: Balsam pear (bitter melon), Chinese waxgourd (Chinese preserving melon), Citrus melon, cucumber, Giberkin, Edible gourds, Melons (including hybrids such as cantaloupe, casaba, crenshaw, honeydew, honey balls, mango melon, muskmelon, and persian melon, pumpkin, summer & winter squash, watermelon.	Aphids, cabbage loopers, cucumber beetles (adults), cutworms, leafhoppers, leafminers, melonworms, pickleworms, plant bugs and squash bugs	Spray plants as needed to ensure uniform coverage. Do not make more than 8 applications per crop. Can be applied up to the day of harvest.
Eggplant	Colorado potato beetles, vegetable leafminers	Spray plants as needed to ensure uniform coverage. Do not make more than 16 applications. Can be applied up to 3 days prior to harvest.
Horseradish	Imported crucifer weevils	Spray as needed. Do not make more than 3 foliar applications. Can be applied up to 22 days prior to harvest.

✓ Leafy Vegetables: amaranth, arrugula, celery, celtuce, chervil, corn salad, chrysanthemum (edible-leaved and garland), cress (garden and upland), dandelion, dick, endive, fennel, lettuce (head & leaf), orach, parsley, purslane (garden & winter) rhubarb, spinach, Swiss chard	Alfalfa loopers, aphids, beet armyworms, cabbage loopers, corn earworms, cutworms, European corn borers, fall armyworm, green cloverworms, southern armyworms, tobacco budworms, vegetable leaf miners.	Spray every 5 to 10 days as needed. Do not make more than 7 applications per crop. Can be applied up to 1 day prior to harvest.
Melons - see Cucurbits		
✓ Potatoes	Beet armyworms, Colorado potato beetles, potato leafhoppers, potato aphids, potato tuberworms/ cabbage loopers, cutworms, European corn borers, potato psyllids, tarnished plant bugs.	Spray plants as needed. Do not make more than 12 applications. Can be applied up to 7 days prior to harvest.

✓ Peppers	Vegetable leafminers, cabbage loopers, flea beetles, corn earworms, pepper weevils, European corn borers.	Spray plants as needed. Do not make more than 8 applications. Can be applied up to 3 days prior to harvest.
✓ Sweet Corn	Corn earworms, European corn borers, fall armyworms, southern armyworms.	Spray plants every 5 days and when needed. Do not make more than 6 applications per season. Can be applied up to 1 day prior to harvest.
✓ Tomatoes (Do not spray on cherry tomatoes and varieties yielding tomatoes less than one inch in diameter).	Beet and southern armyworms, tomato pinworms, tomato fruitworms, hornworms, granulate cutworms, Colorado potato beetles, cabbage loopers.	Spray plants when needed for uniform coverage. Do not make more than 6 applications per season. Can be applied up to date of harvest.

HOME OWNER GROWN FRUIT AND NUT TREES:

Dilute before using. Use clean sprayer. Carefully measure and mix 1 fluid ounce of SPI PERMETHIN HOME & GARDEN and 12 fluid ounces of water. Apply dilute mixture to trees until foliage is dripping wet. Make sure to spray the underside of leaves and into dense foliage. Flush sprayer with clean water after each use.

TREES (Fruit or Nut)	TARGET PESTS	REMARKS
✓ Almonds	Navel orange worms, peach twig borers	Apply when insects appear. Do not make more than 2 applications during the period when the outer hull of the almond has started to open. Do not make more than 5 applications per season. Can be applied up to 7 days prior to harvest.
✓ Apples	Plum curculio, red banded leafrollers, rosyapple aphids, spotted tentiform leafminers, tarnished plant bugs, white apple leafhoppers, oblique-banded leafrollers, green fruitworms.	Apply when insects appear. Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fall.
✓ Peaches	Lesser peachtree borers, peach twig borers, plum curculio, tarnished plant bugs, green fruit worms, Oriental fruit moths	Apply when insects appear. Do not apply within 7 days of harvest. Do not make more than 8 applications per season.

G-(Oct. 11, 00)

Pears	Pear psylla, green fruit- worms and codling moths	Apply when needed. Do not apply more than 3 applica- tions per sea- son. Can be applied up to 14 days prior to harvest.
-------	--	---

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following conditions, disclaimer or warranties and limitations of liability.

In case of Medical emergencies or health and safety inquiries or in case of fire, leaking or damaged containers, information may be obtained by calling

For product information call

**Manufactured by
Speer Products, Inc.
4242 B.F. Goodrich Blvd.
Memphis, TN 38118**

#17