

DESCRIPTION

TEK-TROL is a synthetic detergent with broad spectrum kill of gram positive and gram negative microorganisms, provides effective cleaning and disinfecting in one operation, non-flammable and non-volatile in use dilutions, excellent solubility, dilutes in hot, cold, soft or hard water of up to 1000 ppm water hardness.

The TEK-TROL high foaming detergent system has been especially formulated to provide heavy duty cleaning, wetting and penetration of soils and organic matter and easy rinsing of surfaces. Excellent clinging foam is obtained at 1/2 oz. per gallon of water (1:256) to provide prolonged contact time.

TEK-TROL, at 1/2 oz. per gallon of water (1:256) provides residual control of odor causing bacteria in the presence of moisture.

TEK-TROL is recommended for use when used as directed on ceramic and glazed tile surfaces, stainless steel, aluminum, chrome, galvanized metal, glass, and other hard non-porous surfaces such as treated wood, polyethylene, PVC (polyvinyl chloride), polystyrene, vinyl, fiberglass, viton, ethylene propylene, nitrile, acrylic, polyurethane.

TEK-TROL at 1/2 oz. per gallon of water (1:256) is suited for disinfecting and cleaning in all areas of: swine producing facilities including farrowing and nursery units, finishing floors, grow barns, confinement units, and transport vehicles; poultry, turkey, duck, ratite, and avian rearing facilities, including grow-out and breeder barns, laying operations, and hatcheries, including hatchers and setters, evaporative coolers, fogging systems, humidifying systems, live-haul equipment, chick buses, transfer trucks, coop washing, chick boxes, foot pans, trays, feed bins and buggies; calving facilities including calf hutches and pens, freshening pens, and birthing pens; equine facilities including stables, stalls, breeding and foaling sheds and transport vehicles; hospitals, medical research centers, health care facilities, schools, animal care facilities and research centers, laboratories, veterinary clinics, zoos, and kennels, TEK-TROL can also be used as a wall and floor cleaner in poultry and meat processing plants, meat packing plants, frozen food and canning operations.

TEK-TROL is recommended for use in fogging (wet misting) operations as an adjunct sanitizer of room surfaces either preceding or following regular cleaning and disinfecting procedures. Use a dilution of 1/2 ounce per gallon (1:256). Check with a Bio-Tek representative to determine effective fogging schedule for particular use sites. Leave the room when fogging. In well ventilated areas, such as hatcheries, only 10 to 15 minutes is necessary before reentering. If used in a poorly ventilated area, allow at least 2 hours, or air thoroughly, before reentry. Before fogging in food processing areas, all food and food packaging items must be removed or carefully protected. To thoroughly clean and disinfect room surfaces, before or after fogging, scrub with TEK-TROL at 1/2 ounce per gallon of water (1:256). Disinfected food contact surfaces must be rinsed with potable water prior to reuse.

Toxicity tests required by the U.S. Environmental Protection Agency for TEK-TROL DISINFECTANT-CLEANER have been done in conformity with the EPA Good Laboratory Practice Regulations.

Proven effective as a disinfectant by the A.O.A.C. use dilution test method. All dilutions at 1:256. The following organisms tested at 1000 ppm water hardness (as calcium carbonate) and 10% horse serum (as organic soil):

- Alcaligenes faecalis Strain #838
- Alcaligenes faecalis Georgia Strain
- Aspergillus fumigatus ATCC #8285
- Beta streptococcus, ATCC #19615
- Bordetella bronchiseptica, ATCC #10580
- Candida albicans, ATCC #18804
- A.O.A.C. Fungicidal Test
- Enterobacter aerogenes, ATCC #13048
- Enterococci Group D Species
- Escherichia coli, ATCC # 11229
- Klebsiella pneumoniae, ATCC # 4352
- Microsporium canis, ATCC #42872
- A.O.A.C. Fungicidal Test
- Microsporium gypseum, ATCC #52027
- A.O.A.C. Fungicidal Test
- Mycobacterium bovis (BCG)
- A.O.A.C. (Tuberculoicidal Test)
- Mycoplasma synoviae
- Mycoplasma gallisepticum
- Pasteurella anatipestifer
- Pasteurella multocida, ATCC # 6529 (fowl cholera)
- Proteus vulgaris, ATCC # 13315
- Pseudomonas aeruginosa, ATCC # 15442
- Rhodococcus equi, ATCC #6939

- Salmonella arizonae PD38
- Salmonella arizonae # 7:1, 7, 8
- Salmonella choleraesuis, ATCC #10708
- Salmonella choleraesuis, Kunzendorf Strain
- Salmonella enteritidis, ATCC #13076
- Salmonella gallinarum, ATCC # 9184
- Salmonella pullorum, ATCC # 19945
- Salmonella schottmuelleri, ATCC # 8749
- Salmonella typhimurium, ATCC #13311
- Shigella sonnei, ATCC # 25931
- Staphylococcus aureus ATCC # 6538
- Staphylococcus epidermis
- Staphylococcus hyicus
- Streptococcus equi, ATCC # 39506
- Streptococcus pyogenes, ATCC # 9342
- Streptococcus pyogenes, ATCC # 19615
- Streptococcus suis, Type II
- Taylorella equigenitalis (C.E.M.) Kentucky Strain #3066
- Trichophyton equinum, ATCC #64918
- A.O.A.C. Fungicidal Test
- Trichophyton mentagrophytes, ATCC #9633
- A.O.A.C. Fungicidal Test

*The following viruses tested in accordance with U.S. Environmental Protection Agency Pesticide Assessment Guidelines on hard, non-porous inanimate environmental surfaces. Tested at 1000 ppm water hardness (as calcium carbonate) and 10% horse serum (as organic soil):

- Avian adenovirus
- Avian infectious bronchitis virus,
- Avian influenza virus
- Avian Laryngotracheitis
- Avian reovirus
- Avian rotavirus, ATCC #VR-1
- Canine adenovirus, ATCC #VR-800
- Canine coronavirus, ATCC #VR-809
- Canine parvovirus Strain CPV MLV (Cornell)
- Duck enteritis virus (D.V.E.)
- Equine arteritis, ATCC #VR-796 (Equine viral arteritis)
- Equine herpes virus, ATCC #VR-700 (Equine rhinopneumonitis)
- Equine Rotavirus

- Feline calicivirus, ATCC #VR-782
- Feline rhinotracheitis, ATCC VR-826
- Herpes simplex Type 1
- HIV-1 (AIDS Virus)
- Influenza A-2/Hong Kong virus, ATCC #VR-544
- Mouse hepatitis virus, Strain MHV-A59, ATCC #VR-764
- Newcastle disease virus
- Parainfluenza virus, Type I, Sendai Strain, ATCC #VR-105
- Porcine rotavirus, ATCC #VR-893
- Pseudorabies virus, Aujeszky Strain, ATCC #VR-135
- Transmissible gastroenteritis virus, Purdue Strains, ATCC #VR-763

Hard Surface Mildewicidal Test Method: CSMA method 23 at 1000 ppm water hardness (as calcium carbonate) and 10% horse serum (as organic soil):

- Aspergillus niger, ATCC #6275 (black mold)

- Penicillium variable, NRRL #3765 (green mold)


TEK-TROL

DISINFECTANT • CLEANER CONCENTRATE

FOR USE IN POULTRY, SWINE, EQUINE, VEAL, AVIAN, VETERINARY, RATITE, CALVING, AND ANIMAL CARE FACILITIES

STAPHYLOCIDAL • PSEUDOMONACIDAL • SALMONELLACIDAL
ASPERGILLOCIDAL • TUBERCULOCIDAL • VIRUCIDAL +
FUNGICIDAL • PARVOCIDAL • MILDEWCIDAL • DEODORANT

ACTIVE INGREDIENTS:	ortho-Phenylphenol	12.0%
	ortho-Benzyl-para-Chlorophenol	10.0%
	para-tertiary-Amylphenol	4.0%
INERT INGREDIENTS:		74.0%
	TOTAL	100.0%



EPA EST. NO. 11725-GA-1

KEEP OUT OF REACH OF CHILDREN
DANGER
PRECAUTIONARY STATEMENTS:
Hazard to Humans and Domestic Animals

Danger. Corrosive. Causes eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. May be harmful or fatal if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

Statement of Practical Treatment

IF ON SKIN - Wash with plenty of soap and water. Get medical attention.

IF IN EYES - Hold eyelids open and flush eyes with plenty of water. Get medical attention.

IF SWALLOWED - Call a physician immediately. Drink large quantities of water. Never give anything by mouth to an unconscious person. Avoid Alcohol.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

• CONDUCTIVE •

TEK-TROL has been tested by Hood Patterson-Dewar, Electrical Testing Engineers, for its effect on conductive flooring. The method used conforms to the specifications known as NFPA Standard No. 56A, published by the National Fire Protection Association, Second Edition, 1987. TEK-TROL meets the requirement of NFPA 56A.

Manufactured by:



BIO-TEK INDUSTRIES, Inc.
1380 West Marietta St., NW Atlanta, Ga 30318, USA
Telephone (404) 799-2050 Telex: 671-2844
Fax: (404) 799-2056 Fax: (404) 799-2056

TEK-TROL is a registered trademark of Bio-Tek Industries, Inc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
HOSPITAL USE DIRECTIONS

- To clean, disinfect, and deodorize walls and other hard non-porous surfaces, use 1/2 ounce of TEK-TROL per gallon of water (1:256). Apply with cloth, sponge, mop, mechanical spray device or foaming apparatus, making sure all surfaces are wetted thoroughly. Allow surface to remain wet for 10 minutes. For heavily soiled areas a preliminary cleaning is required; remove gross soil and proceed per above.
- When completed, rinse with potable water.
- TEK-TROL is an effective tuberculocide against Mycobacterium bovis in 10 minutes at 20°C, when used on hard, non-porous inanimate surfaces. Follow above directions, including removal of gross filth or heavy soil prior to application of TEK-TROL.
- KILLS HIV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS.** In health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS),
SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.
— **PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use protective equipment such as disposable rubber gloves, masks or eye coverings.
— **CLEANING PROCEDURE:** Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfectant.
— **CONTACT TIME:** To kill HIV-1 (AIDS virus) a contact time of 10 minutes at room temperature must be allowed.
— **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

HATCHING EGG SANITIZER

TEK-TROL is recommended for use as a Hatching Egg Sanitizer, with best results being achieved in water temperatures ranging from 78° F to 110° F. The recommended dilution is 1/2 oz. per gallon of water (1:256), and may be applied through automatic washing systems, immersion tanks, foaming apparatus, low pressure sprayers, and fogging (wet misting) systems.

GENERAL USE DIRECTIONS

- To clean, disinfect and deodorize walls, floors, tables, drinking fountains, sinks, refrigerators, stoves, restroom fixtures, and garbage cans: Remove gross filth and heavy soil with a preliminary cleaning step prior to application of TEK-TROL. Apply solution of 1/2 ounce of TEK-TROL per gallon of water (1:256) with a sponge, mop, mechanical spray device or foaming apparatus, making sure that all surfaces are wetted thoroughly. Allow surface to remain wet for 10 minutes. Air dry.
- To clean and disinfect such articles as food and water bowls, non-porous bird toys, non-porous perches, combs, brushes, razors, scissors, and rubber goods; all articles must be thoroughly cleaned to remove excess debris, rinsed and rough dried, including the hollows and lumens, before immersion in solution. Soak for 10 minutes in a solution containing 1/2 oz. TEK-TROL per gallon of water (1:256). Rinse food and water bowls prior to reuse.
- To disinfect fabrics such as sheets, linens, aprons and uniforms; First rinse to remove gross filth or heavy soil, then soak for 10 minutes in a solution containing 1/2 oz. of TEK-TROL per gallon of water (1:256).
- To clean gross filth or heavy soil on surfaces prior to disinfection, use 1/2 oz. of TEK-TROL per gallon of water (1:256).
- Dilution ratios: use 1/2 oz. (15cc) of TEK-TROL per gallon of water (1:256) (use 1/16 oz. (2 cc) of TEK-TROL per 16 oz. of water (1:256); 1/8 oz. (4 cc) of TEK-TROL per 32 oz. of water (1:256); 3/16 oz. (6 cc) of TEK-TROL per 48 oz. of water (1:256).

BIO-TEK INDUSTRIES, INC., recommends the use of the BIO-TEK proportioner to provide the automatic method to obtain the 1/2 oz. per gallon solution (1:256).

DIRECTIONS FOR USE IN POULTRY HOUSES, RATITE RANCHES, AVIARIES, FARM PREMISES, SWINE-PRODUCING FACILITIES, EQUINE FACILITIES, VEAL BARN AND CALVING OPERATIONS.

DO NOT USE IN MILKING STALLS, MILKING PARLORS, OR MILK HOUSES.

- Remove all animals and feed from the premises.
- Remove all trucks, coops, crates, water troughs, and feed racks.
- Remove all gross soil, litter, manure, and animal/poultry droppings from floors, walls and surfaces of bams, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals/poultry.
- Thoroughly clean and saturate all surfaces with a solution of 1/2 oz. of TEK-TROL per gallon of water (1:256). Use high pressure sprayer or apply layer of foam with BIO-TEK foaming apparatus.
- Allow to remain in contact for at least 10 minutes
- Immerse halters, ropes and other types of equipment used in handling and restraining animals, forks, shovels, scrapers used for removing litter and manure, in a solution of TEK-TROL at 1/2 oz. per gallon of water (1:256). Allow to remain in contact for 10 minutes.
- Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, waterers, and mangers with a solution of TEK-TROL at 1/2 oz. per gallon of water (1:256). Rinse with potable water before reuse.
- Ventilate buildings, coops and other closed spaces. Do not house poultry or livestock or employ equipment until treatment has been absorbed, set or dried.

BIRD CAGES:

REMOVE BIRDS AND FOOD/WATER BOWLS FROM CAGE.

To clean, disinfect and deodorize bird cages and cage floors, remove gross filth and bird droppings with a preliminary cleaning step prior to application of TEK-TROL. To mix solution of TEK-TROL using calibrated dropper, mix 2 cc of TEK-TROL to 16 oz. of water in the TEK-TROL spray bottle or any other clean spray bottle. For larger solutions, mix 1/2 oz. (15 cc) per gallon of water (1:256). Spray or wipe making sure all surfaces are wetted thoroughly. Allow surface to remain wet for 10 minutes. Air dry. Return birds to cage.

STORAGE AND DISPOSAL

Pesticide Disposal - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
Do not contaminate water, food, or feed by storage or disposal.
Container Disposal - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or purchase and dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NET CONTENTS _____ U.S. GALLONS