

WORN BOLT 2,4-D ESTER

110.3-3

In Accordance with EPA Reg. 100.100-208
NOT REVIEWED
2/25/88

OTHER PRECAUTIONS

To avoid injury to aquatic plants, do not handle or apply the agricultural chemicals with the same equipment used for herbicide 2,4-D. To Vol Ester except as specified on this label.

Do not use 2,4-D herbicide 4 lb. Lo Vol Ester herbicide directly to, or otherwise permit it to come into contact with, corn, grapes, fruit trees, vegetables, flowers or other desirable crop or ornamental plants which are sensitive to 2,4-D herbicide. Do not permit spray mist containing it to drift onto them. Spray only in very small quantities of the spray when it may not be visible. Can cause severe injury during bloom, flowering and dormant periods. Use coarse nozzles to minimize drift. With ground equipment, spray drift can be lessened by keeping the spray boom as low as possible, by applying 20 more gallons of spray per acre, by using no more than 20 pounds spraying pressure with flat fan or fogging flat fan nozzle tips, by spraying when wind velocity is low, and by stopping all spraying when wind exceeds 6 to 7 miles per hour. Do not apply with hollow cone type insecticide or other nozzle that produces a fine droplet spray. With aircraft applications, drift can be lessened by applying not less than 10 gallons of spray per acre, by using no more than 20 pounds spray pressure at the nozzle, by using nozzles which produce a coarse spray, and by spraying only when the wind velocity is not less than 4 miles per hour.

Apply 2,4-D herbicide 4 lb. Lo Vol Ester herbicide only when there is no rain or heavy fog. Do not spray in the vicinity of flowers, buds, tubers or other desirable 2,4-D sensitive crop or ornamental vegetation. Do not spray when the wind is blowing towards susceptible areas of ornamental plants.

At high temperatures (above 95°F) avoid application of this herbicide to susceptible plants growing nearby. Do not use in a greenhouse. Excessive application of this herbicide in the soil may temporarily inhibit seed germination or plant growth.

Local conditions may affect the use of herbicides. Contact your State Agricultural Experiment Station or Extension Service Weed Specialists for advice in selecting treatments from this label to best fit local conditions. Be sure that use of this product conforms to all applicable regulations. Apply this product only as specified on this label.

ENDANGERED SPECIES RESTRICTIONS

The following restrictions apply to use of this product after February 1, 1996:
Before using this pesticide on range and/or pastureland, contact your State Agricultural Experiment Station or Extension Service Weed Specialists for advice in selecting treatments from this label to best fit local conditions. Be sure that use of this product conforms to all applicable regulations. Apply this product only as specified on this label.

RESTRICTED USE PESTICIDES: This pesticide is a Restricted Use Pesticide. It is restricted to use by or under the direct supervision of a person who is certified by the State Agricultural Experiment Station or Extension Service Weed Specialists for the use of this pesticide. The following are the States where this pesticide is registered for use:
Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming.

RESTRICTED USE PESTICIDES: This pesticide is a Restricted Use Pesticide. It is restricted to use by or under the direct supervision of a person who is certified by the State Agricultural Experiment Station or Extension Service Weed Specialists for the use of this pesticide. The following are the States where this pesticide is registered for use:
Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming.

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Read full directions carefully. Do not use if you are pregnant or nursing. Avoid breathing dust or spray. Do not eat, drink, or use tobacco products while using this product. Wash hands thoroughly after use. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

STATEMENT OF PRACTICAL TREATMENT

Apply to the soil in the row or broadcast application. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

ENVIRONMENTAL HAZARDS

Do not use in areas where people or animals are present. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH THE LABELING

Apply Cornen 4 to Lo Vol Ester in water or oil solution in warm weather when weeds or brush are actively growing. Application under drought conditions often will give poor results. Use low spray pressure to minimize spray drift. On cropland and along roadsides, do not exceed 20 psi pressure. Apply enough spray volume to provide uniform coverage of weeds and brush, usually 5 to 20 gallons per acre by ground equipment and 3 to 5 gallons by aircraft. Aerial application may be used if desired to improve spray coverage. Generally, the lower dosage rates recommended on this label will be satisfactory for young, succulent growth of sensitive weed species. Higher dosages are needed for older weeds and under conditions where control is more difficult. The higher dosages may be needed for crop uses. Do not mix with oil or other adjuvants unless otherwise recommended on this label. Deep-rooted perennial weeds such as Canada thistle and other hard weeds and many woody plants usually require repeated applications for maximum control. Do not apply Cornen 4 to Lo Vol Ester where spray drift may contaminate highly susceptible crops or other desirable plants or may contaminate water for irrigation or domestic use. Read and follow all Use Precautions given on this label.

Do not store this product in such a manner as to create or through drift expose workers or the general public. The area being treated must be closed by unobstructed persons. Do not enter treated areas without protective clothing and sprays have dried for appropriate dust hazard (if any).

Some states may require more restrictive entry intervals for various crops treated with this product. Consult your State Department of Agriculture for further information. Workers in the field must be given a warning if they are expected to enter a field of area where this product has been applied. Do not enter treated areas without protective clothing and sprays have dried for appropriate dust hazard (if any). Do not enter treated areas without appropriate protective clothing and sprays have dried for appropriate dust hazard (if any). Do not enter treated areas without appropriate protective clothing and sprays have dried for appropriate dust hazard (if any).

Do not use in areas where people or animals are present. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

Do not use in areas where people or animals are present. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

WEED CONTROL IN SMALL GRAINS NOT UNDERSEEDED WITH A LEGUME: Note: Do not use in areas where people or animals are present.

Spring Wheat and Barley Apply 1 to 2 quarts of Cornen 4 to Lo Vol Ester per acre at the boot stage and when weeds are actively growing. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

Water Wheat and Rye Apply 1 to 2 quarts of Cornen 4 to Lo Vol Ester per acre at the boot stage and when weeds are actively growing. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

Spring Seeded Oats Apply 1 to 2 quarts of Cornen 4 to Lo Vol Ester per acre at the boot stage and when weeds are actively growing. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

Fall Seeded Oats (Southern) Grown for Grain Apply 1 to 2 quarts of Cornen 4 to Lo Vol Ester per acre at the boot stage and when weeds are actively growing. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

Preharvest Treatment Apply 1 to 2 quarts of Cornen 4 to Lo Vol Ester per acre at the boot stage and when weeds are actively growing. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

Preharvest Corn Treatment Apply 1 to 2 quarts of Cornen 4 to Lo Vol Ester per acre at the boot stage and when weeds are actively growing. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

WEED CONTROL IN SORGHUM (MILO) Apply 1 to 2 quarts of Cornen 4 to Lo Vol Ester per acre when sorghum is 6 to 10 inches tall. A high spray pressure may be needed to control some weeds, but the chance of injury to the crop is reduced. Do not use with oil. Do not use with oil. Do not use with oil.

WEED CONTROL IN GRASS SEED CROPS Apply 1 to 2 quarts of Cornen 4 to Lo Vol Ester per acre in the amaranth water required for seed production. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

WEED AND BRUSH CONTROL IN RANGELAND AND GRASS PASTURES Use 2 quarts of Cornen 4 to Lo Vol Ester per acre in water or oil solution. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

WOODY PLANT CONTROL IN NON-CROP AREAS

Apply 1 to 2 quarts of Cornen 4 to Lo Vol Ester per acre in water or oil solution. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

WEED CONTROL IN NON-CROP AREAS OTHER THAN GRASS Apply 1 to 2 quarts of Cornen 4 to Lo Vol Ester per acre in water or oil solution. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

SPRAY TREATMENT To control broadleaf weeds, apply 1 to 2 quarts of Cornen 4 to Lo Vol Ester in water or oil solution. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

Do not contaminate water, food or feeds, drugs or clothing.

This product may be stored in unopened containers in a cool, dry place. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

Do not contaminate water, food or feeds, drugs or clothing.

This product may be stored in unopened containers in a cool, dry place. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

Do not contaminate water, food or feeds, drugs or clothing.

This product may be stored in unopened containers in a cool, dry place. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

Do not contaminate water, food or feeds, drugs or clothing.

This product may be stored in unopened containers in a cool, dry place. Do not use in areas where people or animals are present. Do not use in areas where people or animals are present.

Do not contaminate water, food or feeds, drugs or clothing.

BEST AVAILABLE COPY



CORNBELT[®] 4 LB. LOVOL[®] ESTER

EMULSIFIABLE
LIQUID HERBICIDE

For the control of many broadleaf weeds
and woody plants susceptible to 2,4-D

ACTIVE INGREDIENT		
2,4-Dichlorophenoxyacetic acid		66.0%
Isococyl esters*		
INERT INGREDIENTS		34.0%
		100.0%
2,4-D Dichlorophenoxyacetic Acid Equivalent* 46.95 -- 3.8 lb/gal		
*Isomer Specific by AOAC Method No. 6, DC1-E		

EPA Reg. No. 11773-3

EPA Est. No. 11773 IA-1

Net Contents 2½ Gallons

PRECAUTIONS
Avoid breathing vapors or spray
Wash thoroughly before eating

Statement
In case of contact, wash skin
15 minutes and get medical
attention if the irritant or give full name
person. Get medical attention.

ENVIRONMENTAL
This product is toxic to fish
cleaning of equipment or do
not use for irrigation or drinking
Do not apply this product to
water to protect water.

**IT IS A VIOLATION OF FEDERAL LAW
TO USE THIS PRODUCT IN A MANNER
NOT RECOMMENDED BY THE LABELING**

Apply Cornbelt 4 lb. Lo Vol Ester when
weeds are actively growing. Application
pressure to minimize spray
pressure. Apply enough spray to
5 to 20 gallons per acre by ground
may be used if desired to improve
on this label will be satisfactory
sensitive species and under conditions
needed. For crop uses, do not use
on this label. Deep-rooted perennial
woody plants usually require 2 to
4 lb. Lo Vol Ester where spray or
plants or may contaminate water.
This is given on this label.

Do not use this product in or near
water. The label is being treated.
Do not use this product in or near
water.

Do not use this product in or near
water. The label is being treated.
Do not use this product in or near
water.

Do not use this product in or near
water. The label is being treated.
Do not use this product in or near
water.

Do not use this product in or near
water. The label is being treated.
Do not use this product in or near
water.

Note: If there are uncertainties
regarding use of 2,4-D, consult your
Concise for advice.

To Prepare the Spray: (1) Fill the
sprayer with water. (2) Add
4 lb. Lo Vol Ester in water forms a
suspension. (3) If desired, add
adjuvants to the water. (4) Add
4 lb. Lo Vol Ester to the
mixture. Do not allow
the mixture to become
too thick.

