

PM 23 11773-4 10/11
VAN DIEST SUPPLY CO.

SERVING AGRICULTURE SINCE 1968

HIGHWAY 20 WEST
BOX 610
WEBSTER CITY, IOWA 50595-0610

PHONE 515-832-2366
PLACE ORDERS 1-800-779-2424
FAX 515-832-2955

To: Joanne I. Miller, Product Manager 23
U.S. Environmental Protection Agency
Office of Pesticide Programs
Registration Division (H7505C)
401 M Street, S.W.
Washington, D.C. 20460

CERTIFICATION OF COMPLIANCE

I, being an authorized representative of Van Diest Supply Co., certify that all containers of Cornbelt 6 Lb. LoVol Ester, EPA Registration Number 11773-4; produced by October 23, 1993 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.


I further certify that all containers of said product sold or distributed by this company by April 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.

Tim Nilles
Tim Nilles
Marketing Assistant

2/28/93
Date

Attachment: 2 Copies of Label

Certified: P177 957 590



20711

CORNBELT 6 LB. LOVOL ESTER

EMULSIFIABLE LIQUID HERBICIDE

**Herbicide for the control of many broadleaf weeds,
herbaceous perennials and woody plants
susceptible to 2,4-D in grass pastures,
certain crops and non-crop areas**

ACTIVE INGREDIENT

2,4-Dichlorophenoxyacetic Acid87.7%
Isooctyl esters*

INERT INGREDIENTS 12.3%
100.0%

This product contains 2,4-Dichlorophenoxyacetic Acid Equivalent* 58.3%
-- 5.6 lb./gal.

*Isomer Specific by AOAC Method No. 6.275-6.279 (13th Ed.)

CAUTION

AGRICULTURAL CHEMICAL

KEEP OUT OF REACH OF CHILDREN

See Side Panel For Additional Cautions

EPA Reg. No. 11773-4

EPA Est. No. 11773-IA-1

PRECAUTIONARY STATEMENTS

**HAZARD TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or absorbed through skin. May cause allergic skin reaction. Do not get in eyes, on skin, or on clothing. Avoid breathing vapors or spray mist.

When mixing, loading, or applying this product or repairing or cleaning equipment used with this product, wear eye protection (face shield or safety glasses), chemical-resistant gloves, longsleeved shirt, long pants, socks and shoes. For aerial applicators in enclosed cockpits and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required.

BEST AVAILABLE COPY

Wash hands, face and arms with soap and water as soon as possible after mixing, loading, or applying this product. Wash hands, face and arms with soap and water before eating, smoking, or drinking. Wash hands and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing, loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower.

AGRICULTURAL USE REQUIREMENT: Do not enter or allow agricultural workers to enter into treated areas during the restricted entry interval (REI) of 12 hours.

Note: (1) For containers over 1 gallon and less than 5 gallons: persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron. (2) For containers of 5 gallons or more: A mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Statement of Practical Treatment

In case of contact wash skin with soap and water: for eyes flush with water for at least 15 minutes and get medical attention. If swallowed, DO NOT induce vomiting. Get medical attention. Contains petroleum distillate. NOTE TO PHYSICIAN: May cause chemical pneumonitis if aspirated. If lavage is performed, suggest endotracheal and /or esophagosopic control. Never give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS

Do not contaminate water by disposal of wastes or washing of spray equipment.

This product is toxic to fish. Do not apply directly to water. Apply the product only as specified on the label. Do not contaminate water used for irrigation or domestic purposes.

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.



Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not ship or store with food, feeds, drugs or clothing.

STORAGE

This product may be stored in unheated buildings. Note: Cornbelt 6 lb. LoVol Ester exposed to subfreezing temperatures should be warmed to at least 40 degrees F and mixed thoroughly before use.

DISPOSAL

Pesticide: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Container: PLASTIC: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. METAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Apply Cornbelt 6 lb. LoVol Ester as water or oil spray during warm weather when weeds or brush are actively growing. Application under drought conditions often will give poor results. Use low spray pressure to minimize spray drift. On cropland and along roadsides, do not exceed 20 psi pressure. Apply enough spray volume to provide uniform coverage of weeds and brush, usually 5 to 20 gallons per acre by ground equipment and 1 to 5 gallons by aircraft. Higher gallonage may be used if desired to improve spray coverage. Generally, the lower dosages recommended on this label will be satisfactory for young, succulent growth of sensitive weed species and under conditions where control is more difficult, the higher dosages will be needed. For crop uses, do not mix with oil or other adjuvants unless specifically recommended on this label. Deep-rooted perennial weeds such as Canada thistle and field bindweed and many woody plants usually require repeated applications for maximum control. Do not apply Cornbelt 6 lb. LoVol Ester where spray drift may contact nearby susceptible crops or other desirable plants or may contaminate water for irrigation or domestic use. Read and follow all Use Precautions given on this label.

BEST AVAILABLE COPY

50911

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure). When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "CAUTION Area treated with Cornbelt 6 lb. LoVol Ester on (date of application). Do not enter without appropriate protective clothing until product has dried. In case of accidental exposure wash skin with soap and water; for eyes flush with water for at least 15 minutes and get medical attention. Wash clothing before reuse."

DO NOT USE THIS PRODUCT ON CROPS GROWN FOR SEED. SOME INBREDS HAVE A VERY LOW TOLERANCE TO CHEMICALS AND THE USE OF THIS COMPOUND COULD RESULT IN INJURY TO A CROP GROWN FOR SEED.

Note: If there are uncertainties concerning special local use situations or specific crop variety tolerances to 2,4-D, consult your Agricultural Experiment Station or Extension Service Weed Specialists for advice.

To Prepare the Spray: (1) Fill the spray tank about half full with water then add the required amount of Cornbelt 6 lb. LoVol Ester with agitation, and finally the rest of the water. Note: Cornbelt 6 lb. LoVol Ester in water forms an emulsion which tends to separate unless the mixture is kept agitated. (2) If oil is added, first mix the Cornbelt 6 lb. LoVol Ester and the oil and then add this mixture to the water. However, with adequate agitation, the oil can be added after the Cornbelt 6 lb. LoVol Ester is mixed in the water. (3) If straight oil is used, a solution is formed and separation does not occur. Do not allow any water to get into the oil-herbicide mixture to avoid formation of an invert emulsion.

Local conditions may affect the use of herbicides. For advice in selecting treatments from this label to best fit local conditions, or if there are uncertainties concerning special local use situations or specific crop variety tolerances to 2,4-D, consult local Extension Service or University Specialists for advice.

BEST AVAILABLE COPY

USE PRECAUTIONS


Do not apply Cornbelt 6 lb. Lo Vol Ester herbicide directly to, or otherwise permit it to come into contact with cotton, grapes, fruit trees, vegetables, flowers or other desirable crop or ornamental plants which are sensitive to 2,4-D herbicide. Do not permit spray mist containing it to drift onto them, since even very small quantities of the spray, which may not be visible, can cause injury during both growing and dormant periods. Use coarse sprays to minimize drift. With ground equipment, spray drift can be lessened by keeping the spray boom as low as possible; by applying 20 gallons or more of spray per acre; by using no more than 20 pounds spraying pressure with flat fan or flooding flat fan nozzle tips; by spraying when wind velocity is low; and by stopping all spraying when wind exceeds 6 to 7 miles per hour. Do not apply with hollow cone-type insecticide or other nozzles that produce a fine droplet spray.

With aircraft application, drift can be lessened by applying not less than 5 gallons of spray per acre; by using no more than 20 pounds spray pressure at the nozzles; by using nozzles which produce a coarse spray pattern; and by spraying only when the wind velocity is less than 5 miles per hour.

Applications by aircraft, ground rig and hand dispenser should be carried out only when there is no hazard from spray drift. Do not apply in the vicinity of cotton, grapes, tomatoes or other desirable 2,4-D susceptible crop or ornamental vegetation. Do not spray when the wind is blowing towards susceptible crops or ornamental plants.

At high temperatures (above 95 degrees F) vapors from this product may injure susceptible plants growing nearby. Do not use in a greenhouse. Excessive amounts of this herbicide in the soil may temporarily inhibit seed germination or plant growth.

Do not contaminate irrigation ditches or water used for irrigation or domestic purposes. To avoid injury to desirable plants, do not handle or apply other agricultural chemicals with the same equipment used for Cornbelt 6 lb. LoVol Ester except as specified on this label.



WEEDS CONTROLLED

Cornbelt 6 lb. LoVol Ester herbicide is recommended for control of numerous broadleaf weeds and certain 2,4-D susceptible woody plants without injury to most established grasses. Species controlled include the following, plus many others.

beggarticks	elderberry	pennycress	sowthistle, annual
bitterweed	fanweed	pepperweed, field	spanishneedles
blueweed, Texas	galinsoga	pigweed	sumac
broomweed	garlic, wild	plantains	sunflower
buckbrush	goatsbeard	poorjoe	sweetclover
buckwheat, wild	halogeton	rabbitbrush	tansymustard
burdock	hemp, wild	raddish, wild	tansyragwort
burhead	jewelweed	ragweed	thistle, bull
carpetweed	jimsonweed	rape, wild	thistle, musk
catnip	kochia	redstem	thistle, Russian
chamise	lambquarter	sage, coastal	tumbleweed
chicory	loco, bigbend	sagebrush, big	velvetleaf
cocklebur	mallow, Venice	sagebrush, sand	vervains
coffeeweed	manzanita	salsify	vetch
cornflower	marshelder	sand shinnery	water plantain
coyotebrush	milkvetch	oak	willow
croton	morningglory,	shepherdspurse	witchweed
dandelion	annual	sicklepod	wormwood
docks	mustards	smartweed	yellow rocket
dogfennel	nettles	sneezeweed,	yellow
	onion, wild	bitter	starthistle

WEED CONTROL IN SMALL GRAINS NOT UNDERSEEDED WITH A LEGUME: Note: Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 2 weeks after treatment.

Spring Wheat and Barley: Apply 1/3 to 2/3 pint per acre. Spray when grain is in full tiller stage (usually 4 to 8 inches tall) but before the boot stage and when weeds are small. Do not apply before the tiller stage nor from early boot to the dough stage. Higher rates, up to 1 1/3 pints per acre, may be needed to handle difficult weed problems in certain areas such as under dry conditions especially in western areas. However, do not use unless possible crop injury will be acceptable.

Winter Wheat and Rye: Apply 1/3 to 1/2 pint per acre in the spring at the full tiller stage but before the early boot stage.

Spring Seeded Oats: Apply 1/3 pint per acre at the full tiller stage but before the early boot stage. Oats are less tolerant to 2,4-D than wheat or barley and are more likely to suffer some injury.

Fall Seeded Oats (Southern) Grown for Grain: Apply 1/2 to 1 pint per acre after full tillering but before the early boot stage. Some difficult weeds may require higher rates for maximum control but crop injury may result. Do not spray during or immediately following cold weather.

BEST AVAILABLE COPY

Preharvest Treatment: Apply 2/3 to 1 1/3 pints per acre when grains are in the hard dough stage to control large weeds that may interfere with harvest. Best results will be obtained when soil moisture is sufficient to cause succulent weed growth. **Note:** Do not feed treated straw to livestock.

WEED CONTROL IN CORN: Use one of the following three programs: **Pre-emergence:** Apply 1 1/3 to 2 3/4 pints per acre to soil anytime after planting but before corn emerges. Only emerged broadleaf weeds are likely to be controlled. Do not apply more than 1 1/3 pints per acre unless the increased risk of crop injury can be tolerated. Do not use on light sandy soil. **Emergence:** Apply 2/3 pint per acre just as corn plants are breaking ground. **Postemergence:** After emergence of corn, use 1/3 pint per acre. Application of 1/2 to 2/3 pint per acre may be needed for maximum control of some weeds but such rates are more likely to injure the corn. If the corn is over 8 inches tall, use drop nozzles to keep the spray off the corn foliage as much as possible. Do not apply from 7 to 10 days before tasseling to dough stage. Do not use with oil, atrazine or other adjuvants. Crop injury is more likely to occur if corn is growing rapidly under high temperature and high soil moisture conditions. To reduce breakage of stalks from temporary brittleness caused by 2,4-D, delay cultivation for 8 to 10 days after treatment. Do not forage or feed corn fodder for 7 days following application.

Preharvest Treatment: After the hard dough or denting stage, apply 2/3 to 1 1/3 pints per acre, by air or ground equipment, to suppress perennial weeds, decrease weed seed production and control tall weeds such as bindweed, cocklebur, dogbane, jimsonweed, ragweed, sunflower, velvetleaf and vines that interfere with harvesting. Do not forage or feed corn fodder for 7 days following application.

Note: Hybrids vary in response to 2,4-D and some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Contact seed company, Agricultural Experiment Station or Extension Service Weed Specialists for this information.

WEED CONTROL IN SORGHUM (MILO): Apply 1/3 pint per acre when sorghum is 5 to 15 inches tall. A higher rate of 1/2 to 2/3 pint per acre may be needed to control some weeds but the chance for crop injury is likewise increased. Do not use with oil. Do not treat before the sorghum is 5 inches tall nor during the boot, flowering or early dough stages. If sorghum is taller than 8 inches, use drop nozzles to keep the spray off the foliage as much as possible. Temporary crop injury may occur under the conditions of high soil moisture and high air temperatures. Varieties vary in tolerance to 2,4-D and some hybrids are quite sensitive. Spray only varieties known to be tolerant to 2,4-D. Contact seed company and Extension Service Weed Specialists for this information.

BEST AVAILABLE COPY

WEED CONTROL IN GRASS SEED CROPS: Use 2/3 to 1 pint per acre in the amount of water required for uniform application by air or ground equipment. Apply to established stands in spring from the tiller to early boot stage. Do not spray in boot stage. New spring seedlings may be treated with the lower rate after the grasses have at least five leaves. Perennial weed regrowth may be treated in the fall.

WEED AND BRUSH CONTROL IN RANGELAND, CONSERVATION RESERVE AND GRASS PASTURES: Note: Do not graze dairy animals on treated areas within 7 days after application. Do not graze livestock on treated area within 3 days of slaughter. Do not cut treated area for hay within 30 days after application. Do not use on bentgrasses, alfalfa, clover or other legumes or on newly seeded pastures. Do not apply after heading begins or when grass is in the boot to milk stage where grass seed production is desired. For Conservation Reserve Land follow all applicable State and Federal regulations. Follow the most severe grazing restrictions imposed either by the pesticide label or by the USDA Acreage Conservation Reserve Program whichever is longest.

Bitterweed, Broomweed, Croton, Docks, Kochia, Marshelder, Muskthistle and other Broadleaf Weeds: Use 2 2/3 pints of Cornbelt 6 lb. LoVol Ester per acre in the amount of water needed for uniform application. If the weeds are young and growing actively, 1 1/3 pints per acre will provide control of some species. Deep-rooted perennial weeds may require repeated treatments in the same year or in subsequent years.

Wild Garlic and Wild Onion: Apply 2 2/3 pints per acre, making three applications (fall-spring-fall or spring-fall-spring) starting in late fall or early spring.

WEED CONTROL IN NEWLY SPRIGGED COASTAL BERMUDAGRASS: Apply 1 1/3 to 2 2/3 pints per acre pre-emergence and/or postemergence.

Sand Shinnery Oak and Sand Sagebrush: On the oak, use 1 1/3 pints in 5 gallons of oil or in 4 gallons of water plus 1 gallon of oil per acre. Apply by aircraft between May 15 and June 15. On the sagebrush, use 1 1/3 pints in 3 gallons of oil per acre and apply by aircraft when foliage is fully expanded and the brush is actively growing.

Big Sagebrush and Rabbitbrush: Use 2 2/3 to 4 pints per acre in 2 to 3 gallons of oil or in 3 to 5 gallons of oil-water emulsion spray. For rabbitbrush, the 4 pint rate is usually required. Brush should be leafed out and growing actively when treated. Retreatment may be needed.

Chamise, Manzanita, Buckbrush, Coastal Sage, Coyotebrush and Certain Other Chaparral Species: Use 2 2/3 to 4 pints per acre in 5 to 10 gallons of water. One gallon of fuel oil may be included in the spray mixture for added effectiveness. Make applications by aircraft or ground equipment to obtain uniform spray coverage. For effective control, the brush must be fully leafed out and growing actively when sprayed. Retreatment may be needed.



Woody Plant Control in Non-Crop Areas: To control species susceptible to 2,4-D in right of ways, fencerows, roadsides, and along drainage ditch banks, spray brush up to 5 to 8 feet tall after spring foliage is well developed, using 4 to 5 pints of Cornbelt 6 lb. LoVol Ester in 100 gallons of water and wetting all parts of the brush including foliage, stems and bark. This may require up to 400 gallons of spray per acre for adequate coverage of solid stands of brush. Make application in such a way as to prevent drift of the spray off the area being treated. Spraying can be effective at any time up to 3 weeks before frost as long as soil moisture is sufficient for active growth of the brush. Control will be less effective in mid-summer during hot dry weather when soil moisture is deficient and plants are not actively growing. Oil or wetting agent may be added to the spray, if needed for increased effectiveness.

WEED CONTROL IN NON-CROP AREAS SUCH AS LAWNS, GOLF COURSES, CEMETERIES, PARKS, AIRFIELDS, ROADSIDES, VACANT LOTS, DRAINAGE DITCH BANKS: After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled. Apply 1 1/3 to 2 2/3 pints of Cornbelt 6 lb. LoVol Ester per acre in the amount of water needed for uniform application. Usually 2 2/3 pints per acre provide good weed control under average conditions. Treat when weeds are young and growing well. Do not use on golf greens nor on dichondra or other broadleaf herbaceous ground cover. Do not use on creeping grasses such as bent and St. Augustine except for spot treating, nor on newly seeded turf until grass is well established. Reseeding of treated areas should be delayed following treatment. With spring application, reseed in the fall; with fall application, reseed in the spring. Legumes are usually damaged or killed so do not treat areas where legumes are desired. Deep-rooted perennial weeds may require repeated treatments. **RESTRICTIONS ON RETREATMENT:** The maximum number of broadcast applications per treatment site is 2 per year.

Tule (Bulrush) and Other Rushes: Mix 2 2/3 pints of Cornbelt 6 lb. LoVol Ester and 1 gallon of diesel oil or kerosene, then add this mixture to 100 gallons of water. Spray to wet all foliage (400-800 gallons per acre). Addition of a wetting agent may be advisable. Apply in the spring during flower head emergence. Respray if needed when regrowth is 3 to 5 feet tall.

SPOT TREATMENT: To control broadleaf weeds in small non-cropland areas with a hand sprayer, use 1/6 pint of Cornbelt 6 lb. LoVol Ester in 3 gallons of water and spray to thoroughly wet all weed foliage. Keep spray mixture agitated to prevent separation.



WARRANTY STATEMENT

1. Van Diest Supply Co. warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Van Diest Supply Co. neither makes, nor authorizes any agent or representative to make any other warranty of Fitness or of Merchantability, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Van Diest Supply Co.'s control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Van Diest Supply Co. under 1 above) resulting from handling, storage, and use of this material.

Cornbelt is a registered trademark of Van Diest Supply Co., Webster City, Iowa 50595.

**Manufactured by Van Diest Supply Co.
Webster City, Iowa 50595**

