



011773-00016-052599

Printed: 09:44:45 Thursday, 24 Jun, 1999 # 29 / 3429

Systems Integration Group, Inc.

PM23

11773-16

5/25/99

1 of 4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

MAY 25 1999

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Tim Niles
Van Diest Supply Company
1434 220th Street
P.O. Box 610
Webster City, IA 50595

Dear Mr. Niles:

Subject: Label Amendment Extending 2,4-D Use in Reduced or No-Tillage Soybeans
(Pre-plant Only)
Cornbelt Salvan
EPA Reg. No. 11773-16
Your Application Dated April 24, 1999

The Agency is conditionally approving an amendment to the registration of the above-referenced product under the authority of section 3(c)(7)(B) of the Federal Insecticide, Fungicide Act (FIFRA). This amendment allows use of the subject product on reduced or no-tillage soybeans (pre-plant only) with a maximum permissible level for residues of the herbicide in or on soybeans of 0.1 ppm. This amendment will expire automatically on December 31, 2001. In addition, during the period that this amendment is effective, it will be subject to the conditions listed below:

- 1) This acceptance is based on your certification that the submitted labeling on preplant soybean use is an exact copy of the labeling accepted by the Agency on September 25, 1996. Please note that the acceptance of this labeling pertains to the previously approved preplant soybean use only and does not signify the acceptance of any other labeling revisions. No other label revisions are being reviewed or considered with this action. You should also note that if you fail to satisfy the conditions imposed in this certification, EPA may issue a notice to cancel this amendment under FIFRA section 6(e).
- 2) This conditional registration will expire automatically on December 31, 2001. Sale or distribution of the subject product bearing labeling for this use on reduced or no-tillage soybeans (pre-plant only) after December 31, 2001 will be illegal. The tolerance authorizing residues of the subject product will also expire on December 31, 2001. After that date, sale or distribution of food in interstate commerce containing any residue of the subject product will be a violation of the Federal Food, Drug, and Cosmetic Act.

M Coakle
5-25-99

- 3) Finally, if and when a permanent tolerance is established, EPA will entertain an application to amend the registration of the subject product without any special limitations on the duration of the amendment.

A stamped copy of the labeling is enclosed for your records. Please submit one (1) copy of your final printed labeling before you release the product for shipment.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

3074

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAY 25 1999

VAN DIEST SUPPLY CO.
Webster City, Iowa 50595

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

SUPPLEMENTAL LABELING - CORNBELT SALVAN (EPA Reg. No. 11773-16)

11773-16

For Use In Crop Residue Management Systems in Soybeans
(Preplant Application Only)

ATTENTION

- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Cornbelt Salvan herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Cornbelt Salvan herbicide according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container and this supplemental labeling.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

General Information

Cornbelt Salvan herbicide may be used for postemergence control of many susceptible annual and perennial broadleaf weeds. This product may be applied prior to planting soybeans to provide foliar burn-down control of susceptible annual and perennial broadleaf weeds and certain broadleaf cover crops such as those listed on this label. Make only preplant applications to emerged weeds prior to planting soybeans grown in reduced tillage production systems. Apply only according to instructions given below.

Do not use any tillage operations between herbicide application and planting of soybeans.

Mixing Instructions

Compatible crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may be added to spray mixtures to increase the herbicidal effectiveness of Cornbelt Salvan herbicide on certain weeds. Read and follow all directions and precautions on this label and on the label of each product added to the spray mixture.

Application Procedures

Apply using air or ground equipment in a spray volume sufficient to provide uniform coverage of weeds. Use 2 or more gallons of total spray volume per acre for aerial application and 10 or more gallons per acre for ground equipment.

Application Timing and Use Rates

Product	Broadcast Application Rate	When To Apply (Days Prior to Planting Soybeans)
Cornbelt Salvan	.8 pt/acre 1.6 pts/acre	Not less than 7 days Not less than 30 days

(Continued On Back)

Weeds Controlled

arrowhead	hemp dogbane	puncture vine
artichoke	henbit	purslane
beggar ticks	horsetail	ragweeds
bindweeds	jimsonweed	russian thistle
blue thistle	klamathweed	quickweed
bull thistle	kochia	shepherd's purse
burdock	lambsquarters	smartweed
butterprint	loco weed	sow thistle, annual
canada thistle	mallow	stinkweed
catnip	marestail	sumacs
chickweed	mexican weed	sunflower
chicory	morning glory	wild garlic
cocklebur	musk thistle	wild mustard
dandelion	mustards	wild onion
docks	nettles	wild parsnip
fanweed	pennycress	wild radish
figwort	pepper grass	wild rape
four o'clock	pigweed	wild sweet potato
ground ivy	prickly lettuce	yellow goatsbeard

Some weeds above, listed as susceptible may not be completely killed but growth is so retarded by proper application that they are no longer a factor affecting growth and yield of crop.

For best weed control results, application should be made when weeds are small, actively growing and free of stress caused by temperature extremes, moisture stress, diseases, or insect damage. The control of individual weed species may be variable. Consult your local county agent or state agricultural extension specialist or crop consultant for advice.

Use Precautions and Restrictions

- Important Notice: Unacceptable injury to soybeans planted in treated fields may occur. Whether or not soybean injury occurs and the extent of such injury will depend on weather (temperature and rainfall) from herbicide application until soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present at the time of application. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.
- Do not use on sandy soils with less than 1.0% organic matter.
- Do not make more than one application per season regardless of the application rate used.
- Do not apply when weather conditions such as atmospheric temperature inversion or when wind direction favors drift from the treated area to susceptible plants.
- Do not allow livestock grazing or harvest hay, forage or fodder from treated fields. Livestock must be restricted from feeding/grazing of treated cover crops.
- In treated fields, plant soybean seed as deep as practical, but not less than 1.0 inch deep. Adjust the planter, if necessary, to ensure that planted seed is adequately covered.
- Do not apply Cornbelt Salvan herbicide as described in this supplemental labeling unless you are prepared to accept the results of soybean injury, including possible stand loss and/or yield reduction.
- During the growing season following application, do not replant treated fields with crops other than those labeled for use with Cornbelt Salvan herbicide.