


DE-BUG-1

For Lawn, Garden & Agricultural Control of Grasshoppers, Crickets, Army Worms, Cutworms, Darkling Ground Beetles, Mole Crickets, Earwigs and Sow Bugs

Active Ingredients By Weight:

Carbaryl (1-naphthyl methylcarbamate).....	2.0%
Inert Ingredients.....	98.0%
Total.....	100.0%



KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

N-Methyl Carbamate


- IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat.
- IF ON SKIN: Wash thoroughly with soap and water.
- IF IN EYES: Flush with plenty of water.

NOTICE OF WARRANTY

The manufacturer warrants that the material contained herein conforms to the chemical description on the label for the purpose stated in the Directions for Use. When purchasing the product, the buyer assumes all risks of the handling and use which may result in damage or losses that are beyond the control of the seller; as in the event of incorrect method of application, incompatibility with other products used, or the presence on crop or soil of other substances. There is no other warranty made by the manufacturer as to the fitness or merchantability. The limit of liability of the manufacturer or any seller for any losses, damages or injuries resulting from the use of this product will be the purchase price paid for this product by the buyer or user for the quantity involved. No verbal or written change to this warranty is valid. All buyers and users of this product are understood to have accepted all terms and conditions of this warranty.

EPA EST. NO. 11773-NE-001

EPA REG. NO. 11773-18



### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

**CAUTION:** May be harmful if swallowed. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

**NOTE TO PHYSICIAN:** 1-Naphthyl Methylcarbamate is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

**Personal Protective Equipment:**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENVIRONMENTAL HAZARDS


This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**Pesticide Disposal:** Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by State and local authorities.



### GENERAL INFORMATION


Read entire label. Use in strict accordance with the label directions and cautions.

**GENERAL:** Broadcast application may be made with spreader, hand applicator, aircraft or by hand. Use gloves and wash thoroughly following application. Treatment may be repeated as necessary.

**CROP AND RANGELAND:** Apply 2 to 20 pounds per acre for control of grasshoppers, crickets and cutworms.

**HOME AND GARDEN:** Use ½ to 2 ½ pounds per 10,000 square feet of area. Use the higher application rates when insect density is high and crops or garden are threatened.

**FOR SUPPRESSION** of grasshoppers and crickets, rates of ½ to 1 ½ pounds per acre may be used.



### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for the training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

### IMPORTANT PRE-HARVEST INFORMATION

Tolerances established under the Federal Food, Drug, and Cosmetic Act permit the sale of crops bearing probable carbaryl residues when carbaryl is used in accordance with label directions. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).



## CROP AND DAYS BETWEEN APPLICATION &amp; HARVEST

Almond	28 days
Apples, apricots, broccoli, brussel sprouts, cabbage, cauliflower, cherries, cucumbers, eggplants, melons, okra, peaches, pears, peppers, squash, tomatoes	3 days
Asparagus	1 day
Beans, peas	3 days (succulents) 14 days (forage) 21 days (dried, hay)
Alfalfa, carrots, garden beets (roots), horseradish, parsnips, potatoes, radishes, rutabagas, strawberries, turnips (roots)	7 days
Citrus	5 days
Collard, endive (escarole), forest land, garden beets (tops), grassland, hay, kale, leaf lettuce, parks, parsley, pastures, rangeland, recreation areas, spinach, swiss chard, turnip (tops), walnut, wasteland	14 days
Corn (field and pop)	14 days (forage, silage) 48 days (grain, fodder)
Corn (sweet)	2 days (ears) 14 days (forage) 48 days (fodder)
Head lettuce	14 to 15 days
Sugar beets	28 days (roots, forage)
Wheat	21 days

Van Diest Supply Co.  
P.O. Box 410  
McCook, NE 69001-0410