



DRY DIP INSECTICIDE

A powdered insecticide for use on Livestock
and Poultry. With deodorant.

ACTIVE INGREDIENTS:

Malathion* 5%

INERT INGREDIENTS: 95%

*O,O-dimethyl phosphorodithioate of diethyl
mercaptosuccinate



CAUTION

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed. Avoid breathing of dust. Avoid contact with skin. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Apply this product only as specified on the label. Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of lakes, streams or ponds.

DIRECTIONS

POULTRY: (Northern Fowl Mite, Poultry Lice) Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4 to 8 weeks or when necessary. Treatment of poultry for Northern Fowl Mite should be a supplemental treatment to the treatment of the premises. **PREMISE TREATMENT**—Apply 1 lb. per 50 to 60 sq. ft. of litter and floor space and to nests, roosts and adjacent areas.

HOGS: (Lice) Treat animals thoroughly. Repeat in 10 days and thereafter as needed. Thoroughly dust the pens in addition to dusting the swine.

DOGS and CATS: (Fleas) Direct application: Apply sufficient dust to treat animal thoroughly. Use any garden type duster, puff duster or shaker-type can. If animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animals should be closely examined for fleas between treatments. If the animal is confined to an area free of fleas, one treatment may be sufficient.

EPA Reg. No. 11849-2

CATTLE (Beef and Non-Milking): (Lice) Treat Animal thoroughly. Repeat in 10 days and thereafter as needed. (Horn Fly) Use 3 tbs. per animal. Apply on back and neck of the animal and repeat applications at 10-14 day intervals. **NOTE:** Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening.

DAIRY CATTLE: (Horn Fly) Use 3 tbs. per animal. Apply on back and neck of the animal. Do not apply to dairy cattle later than 5 hours before milking or during milking. Repeat applications at 10-14 day intervals. Do not treat animals under one month of age. When applying dusts avoid contamination of feed, food containers and watering troughs.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

MBSCO

NET WEIGHT 25 LBS.



Manufactured by

SILAK COMPANY

HOSPERS, IOWA 51238

