

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May be harmful if swallowed. Do not breathe dust. Do not get in eyes, on skin, or on clothing. May cause eye and skin irritation. During application, treatment and loading operations, wearing gloves, clean synthetic rubber gloves, and on return to clothing. Wash hands, arms and face thoroughly with soap after handling and before eating or drinking. Routinely shower and change clothes and wash work clothes with soap and water after use.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and other wildlife. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

Use according to label directions for specific pests. Do not use in a manner inconsistent with this label.

STORAGE AND DISPOSAL

PROHIBITION: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Waste containing this product may be hazardous to the environment. Do not dispose of in a manner inconsistent with this label.

CONTAINER DISPOSAL: If not reusable, completely empty container by shaking and tapping sides and bottom to loosen clinging material. Empty residue into application equipment. Then dispose of liner in a sanitary and/or by incineration if allowed by State and local authorities. Do not reuse parts or pieces of the same manner.

STORAGE: Store in a cool, dry place.



DRY DIP INSECTICIDE

A powdered insecticide for use on Livestock and Poultry

ACTIVE INGREDIENTS:

Malathion* 5%

INERT INGREDIENTS: 95%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate



KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes for at least 15 minutes with water. Call a physician if irritation persists.

NOTE TO PHYSICIAN- ANTIDOTE STATEMENT

If signs of cholinesterase inhibition are present atropine is antidote. 2-4 mg is also antidote and may be administered in conjunction with atropine.

See side panel for additional precautionary statements.

EPA Reg No. 11849-2 EPA Est. No. 11849-1A-2

NET WEIGHT 25 LBS.



Manufactured By
InterChem, Inc.
Alton, Iowa 51003

ACCEPTED

DEC 3 1984

Under the Pesticide
Fungicide, and B
in accordance, for the
registered under
EPA Reg. No. 11849-2

10F1
Pm 16
11849-2

DIRECTIONS

FOR USE ON ANIMALS AND POULTRY:

POULTRY: (Northern Fowl Mite, Poultry Lice) Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4 to 6 weeks or when necessary. Treatment of poultry for Northern Fowl Mite should be a supplemental treatment to the treatment of the premises. **PREMISE TREATMENT:** Apply 1 lb. per 50 to 60 sq. ft. of litter and floor space and to nests, roosts and adjacent areas.

HOGS: (Lice) Treat animals thoroughly. Repeat in 10 days and thereafter as needed. Thoroughly dust the pens in addition to dusting the swine.

DOGS AND CATS: (Fleas) Direct application. Apply sufficient dust to treat animal thoroughly. Use any garden type duster, puff duster or shaker type can. If animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animals should be closely examined for fleas between treatments. If the animal is confined to an area free of fleas, one treatment may be sufficient. Do not treat kittens and puppies under 4 weeks old.

CATTLE (Beef and Non-Milking): (Lice) Treat animal thoroughly. Repeat in 10 days and thereafter as needed. (Horn Fly) Use 3 tbs. per animal. Apply on back and neck of the animal and repeat applications at 10-14 day intervals. **NOTE:** Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening.

DAIRY CATTLE: (Horn Fly) Use 3 tbs. per animal. Apply on back and neck of the animal. Do not apply to dairy cattle later than 5 hours before milking or during milking. Repeat applications at 10-14 day intervals. Do not treat animals under one month of age. When applying dusts avoid contamination of feed, food containers and watering troughs.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of misuse or improper handling and accepts the product on these conditions.

HP

LOT NO.

BEST AVAILABLE COPY