

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

May be harmful if swallowed. Do not breathe dust. Do not get in eyes, on skin, or on clothing. May cause eye or skin irritation. During exposure in treatment and loading operations wear goggles, clean synthetic rubber gloves, and protective clothing. Wash hands, arms, and face thoroughly with soap after handling and before eating or smoking. Routinely shower or bathe after work and wash work clothes with soap and hot water before reusing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: (Plastic pail with liner) Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. Do not reuse pail, dispose of in the same manner.

STORAGE: Store in a cool, dry place.



DRY DIP INSECTICIDE

A powdered insecticide for use on Livestock and Poultry

ACTIVE INGREDIENTS:

Malathion: 0,0-dimethyl phosphorodithioate of diethyl mercaptosuccinate

INERT INGREDIENTS:

5%
95%

CONTAINS

CYTHION

THE PREMIER GRADE MALATHION

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED - Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED - Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

IF ON SKIN - Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES - Flush eyes for at least 15 minutes with water. Call a physician if irritation persists.

NOTE TO PHYSICIAN - ANTIDOTE STATEMENT

If signs of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. See side panel for additional precautionary statements.

EPA Reg. No. 11849-2 / EPA Est. No. 11849-IA-2

NET WEIGHT 25 LBS.



Manufactured By

WHB SPECIALTY PRODUCTS CO.
P.O. Box 231 • Alton, Iowa 51003

DIRECTIONS FOR USE ON ANIMALS AND POULTRY:

POULTRY: (Northern Fowl Mite, Poultry Lice) Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4 to 8 weeks or when necessary. Treatment of poultry for Northern Fowl Mite should be a supplemental treatment to the treatment of the premises.
PREMISE TREATMENT: Apply 1 lb. per 50 to 60 sq. ft. of litter and floor space and to nests, roosts and adjacent areas.

HOGS: (Lice) Treat animals thoroughly. Repeat in 10 days and thereafter as needed. Thoroughly dust the pens in addition to treating the swine. Do not treat hogs under 1 month of age.

DOGS AND CATS: (Fleas) Direct application. Apply sufficient dust to treat animal thoroughly. Use any garden type duster, puff duster or shaker-type can. If animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animals should be closely examined for fleas between treatments. If the animal is confined to an area free of fleas, one treatment may be sufficient.

Do not treat kittens and puppies under 4 weeks old.

CATTLE (Beef and Non-Milking): (Lice) Treat animal thoroughly. Repeat in 10 days and thereafter as needed. (Horn Fly) Use 3 lbs. per animal. Apply on back and neck of the animal and repeat applications at 10-14 day intervals. **NOTE:** Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening.

DAIRY CATTLE (Lactating): (Horn Fly) Use 3 lbs. per animal. Apply on back and neck of the animal. Do not apply to dairy cattle later than 5 hours before milking or during milking. Repeat applications at 10 - 14 day intervals.

Do not treat cattle under 3 months of age.

Use only with adequate ventilation. After using this product indoors, ventilate thoroughly before occupying enclosed spaces.

When applying dusts avoid contamination of feed, food containers and watering troughs.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of misuse or improper handling and accepts the product on these conditions.

HP 3357

LOT NO.

BEST AVAILABLE COPY

10F1
PM 16
11849-2

RECEIVED
FEB 27 1989

11849-2