

RAMEX

RAT and MOUSE KILLER

Manufactured exclusively with Toxivil.

CAUTION
KEEP OUT OF REACH
OF CHILDREN

RAMEX contains as its active ingredient an anti coagulant chemical. In case baits are eaten, give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately. Since all effective roderenticides are toxic, every precaution should be exercised in their use.

ACTIVE INGREDIENT


"Diphacinone
(2-Diphenyl-1-picrylhydrazide) 0.005%

INERT INGREDIENTS 99.995%

Do not reuse container. Destroy when empty.

NOTE TO REGISTRARS

If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdose of bishydroxycoumarin.



Do not reuse container. Destroy when empty.

NOTE TO PHYSICIANS

If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdose of bishydroxycoumarin.

WHEN BAITING OUTDOORS, TO PROTECT CHILDREN, PETS, DOMESTIC ANIMALS AND WILDLIFE, USE TAMPER-PROOF BAIT STATIONS ONLY.

DIRECTIONS

(Before using, read directions carefully)

RATS — Suggested Use: Place a minimum of 4 ounces of bait where rats are known to feed, water or travel. An uninterrupted supply of bait should be maintained for approximately 10 days. Bait placements along walls, in corners and in concealed locations where rats and mice can feed unobserved are desirable.

MICE: Place tablespoon amounts of bait at 8-12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days. Baiting should be continued as long as evidence of feeding is observed. Where a continuous source of infestation is present from nearby dumps or fields permanent tamper-proof bait stations should be used and bait placed in them as often as necessary. Contaminated or spoiled bait should be replaced promptly.

NET CONTENTS:



EPA Reg. No. 11849-1

Manufactured by

SILAK COMPANY, INC.
Hospers, Iowa 51238

RAMEX

RAT and MOUSE KILLER

Manufactured exclusively with Toxivil.

RAMEX contains as its active ingredient an anti coagulant chemical. In case baits are eaten, give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately. Since all effective rodenticides are toxic, every precaution should be exercised in their use.

ACTIVE INGREDIENT

If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdose of bishydroxycoumarin.