

DIRECTIONS FOR USE

IMPORTANT: Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using SILAK 804C. For an aid in reducing Face Fly populations, faces of cattle must be dusted daily. Best results are achieved when dust applicators are used as directed on an area wide basis.

CONDITIONS OF SALE: These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. SILAK COMPANY and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accept the product on these conditions.

SILAK 804C

ANIMAL INSECTICIDE

CONTAINS CO-RAL®

For control of horn flies and lice and aids in reducing face fly population, on beef and dairy cattle, and control of lice on swine.

ACTIVE INGREDIENT:

O,O-Diethyl O-(3-chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl) phosphorothioate*

1.00%

INERT INGREDIENTS

99.00%

100.00%

EPA Reg. No. 11849-7

*CO-RAL is a Reg. TM of The German Trademark Proprietor, Leverkusen

NET WT _____ LBS. LOT NO _____

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Side Panel for Additional Cautions.



Manufactured by
SILAK COMPANY, INC.

Hospers, Iowa 51238

RECOMMENDED APPLICATIONS

CATTLE (Dairy and Beef): (Horn Flies & Lice) Direct Application: Apply not more than 2 oz. per animal. Dust evenly into the hair over the head, neck, shoulders, back and tailhead. Repeat as necessary. No interval is required between treatment and use of meat or milk as food.

SWINE: (Lice) Direct Application: Apply not more than 1 oz. per animal as a uniform coat to the shoulders and back. Repeat as necessary, but not more often than once every 10 days. No interval is required between treatment and use of meat as food. **Bedding Treatment:** Apply 2 oz. uniformly over each 30 square feet of fresh, dry bedding. Repeat as necessary, but not more than once every 10 days. **Direct Application and Bedding Treatment:** In severe infestations, both individual animals and the bedding may be treated as directed above.

RESTRICTIONS

For external insecticidal use on above specified animals only. Avoid contamination of feed, troughs, water and water utensils. Provide thorough ventilation while dusting.

CAUTION

Harmful if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate feed or food. Keep out of reach of children. If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

To Physician - Atropine sulfate is antidotal. 2-PAM is also antidotal, and may be administered in conjunction with atropine.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.