

ACCERTED

Oct 17 1973

FONTE
UNIVERSITY HOSPITAL

**CAUTION
KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. Avoid inhalation of mist and contact with skin. In case of contact with skin or eyes flush with plenty of water; for eyes get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. If swallowed induce vomiting immediately and get medical attention.

Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Do not reuse container. Destroy when empty.

DO NOT USE POUR SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Do not treat animals within one day of slaughter.

Do not apply dermally in combination with Trichlorfon.



GENERAL DIRECTIONS

A general insecticide used effectively against horn flies, stable flies, face flies, house flies, and other flying insects. Not to be applied directly to the skin of farm animals. Never apply to pregnant cattle or to calves under 6 months of age. Apply to the power or electrically charged metal stems of the horns, and to the neck and head of the animal. Do not wet the skin of the animal or allow it to rain. Do not apply regularly, as it may cause death. Bees and other insects should not be sprayed as they may show sensitivity to organic phosphates.

**CIODRIN - VAPONA
INSECTICIDE
READY-TO-USE
ANIMAL SPRAY SOLUTION**

A Ciodrin® and Vapona® Insecticide Ready-to-Use Fly Spray For Dairy and Beef Cattle. Controls Horn Flies, Stable Flies, Face Flies, House Flies and Lice.

ACTIVE INGREDIENTS	BY WEIGHT
Dimethyl Phosphate of Alpha-Methyldene- 3-Hydroxy-Cis-Crotonate*	1.00
2,2-Dichlorovinyl Dimethyl Phosphate	0.25**
Related Compounds	0.025*
TOTAL	1.275
PER 100 ml	

INERT INGREDIENTS	PER 100 ml
Water	100

**CAUTION
KEEP OUT OF REACH OF CHILDREN**

See Other Cylinders for Precautions



Manufactured by
SHARK COMPANY, INC.
Hosiers, Iowa 51238

**USE AND DIRECTIONS FOR
APPLICATION**

LACTATING DAIRY AND BEEF ANIMALS: Flies, Horn Fly, Stable Fly, Face Fly, House Fly. For the control of horn flies, stable flies, face flies, and house flies, make a light application of CIODRIN-VAPONA READY-TO-USE INSECTICIDE daily as a fine mist to thoroughly cover all parts of the animal, including the legs.

LACTATING DAIRY AND BEEF ANIMALS: Lice. Make 1-2 applications of 3-3½ ounces of CIODRIN-VAPONA READY-TO-USE INSECTICIDE at 14 day intervals as a fine mist spray to thoroughly cover all parts of the animal. If material is applied by fogging or misting equipment, the maximum spraying time per animal is 30 seconds.

BACK AND FACE RUBBERS: For the control of horn flies and face flies on lactating dairy and beef animals, use CIODRIN-VAPONA READY-TO-USE INSECTICIDE on back rubbers and face rubbers. Use 1 gallon for each 20 linear feet of burlap back rubber. For automatic back rubbers, recharge the reservoir as needed. Animals must rub face on rubbers for best face fly control.

NOTICE: Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on this basis.

EPA Reg. No. 6106

14
RECEIVED
OCT 1 1973
Pesticides Regulation
Division EPA

NET CONTENTS