

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING! HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN.
DO NOT BREATHE SPRAY MIST. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING.
Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water. For eyes, get medical attention. Wash all contaminated clothing with soap and hot water before reuse. Do not breathe fumes. Rapidly absorbed through skin.

During commercial exposure in spray mixing and loading operations, wear clean, natural rubber gloves, waterproof protective clothing and goggles when handling. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part II. Keep all unprotected persons out of operational area.

Precautions must be observed when handling both the concentrated material and the diluted emulsions. Store the concentrate in a safe place. Children or animals must not be able to reach or handle the concentrate or diluted emulsions.

ENVIRONMENTAL HAZARDS

The product is toxic to fish, birds and other wildlife. Keep out of rivers, streams or ponds. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Do not store in or around fire line. Store the concentrate in a safe place. Children or animals must not be able to reach or handle the concentrate or diluted emulsion.

PESTICIDE DISPOSAL: Pesticides are often extremely potent and/or caustic. Do not spray mixture or residue down drain. It is a Federal law that use will be discontinued if used by use according to label directions, contact your State Fertilizer and Pesticide Control Agency or their State and Water Resources Institute at the U.S. EPA Region Office for guidance.

CONTAINER DISPOSAL (METAL): Triple rinse container. Then puncture, cut or recondition, or puncture and dispose of in a sanitary, non-contaminated receptacle approved by state and local authorities.

Decontaminate used containers by washing thoroughly with a strong solution of liquid detergent and rinsing with water. Dispose of rinsings away from surface water and/or contaminate water supplies. **CAUTION:** Avoid skin and eye contact with caustic solutions.

CONTAINER DISPOSAL (PLASTIC): Triple rinse container. Then puncture, cut or recondition, or puncture and dispose of in a sanitary, non-contaminated receptacle. If allowed by state and local authorities, by burning, if burned, stay out of smoke.



CIODRIN VA.PONA INSECTICIDE

Emulsifiable Concentrate

CONTAINS
CIODRIN and VAPONA Insecticides

ACTIVE INGREDIENTS	BY WEIGHT
Dimethyl phosphate of alpha-methylbenzyl	10.0%
3-hydroxy-cis-crotonate	10.0%
2,2-dichlorovinyl dimethyl phosphate	2.3%*
Related Compounds	0.2%
Xylene	77.0%
INERTINGREDIENTS	10.5%
TOTAL	100.0%

* CIODRIN® Insecticide
** Equivalent 2.5% Vapona (DDVP) Insecticide

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENT OF PRACTICAL TREATMENT

IF INHALED: Call physician or poison control center immediately. If irritant presents aspiration hazard, if possible vomit fingers in the mouth. Under medical supervision drink one or two glasses of water and induce vomiting by means of emetics. If breathing difficulty continues, call physician or go in any emergency room. Do not induce vomiting unless directed by physician.

IF IN SKIN: Remove victim to fresh air. Apply normal respiration if breathing stopped.

IF ON SKIN: Remove victim to fresh air. Wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call physician or go in any emergency room.

NOTE TO PHYSICIAN: ANTIDOTE STATEMENT
For insecticide exposures, call physician or go in any emergency room.

See side back panel for additional precautionary statement.

NET CONTENTS (LITERS)

EPA Reg. No. 11E49-9 EPA Est. No. 11E49-IA-2

Manufactured By

InterChem, Inc.
Alton, Iowa 51003

DIRECTIONS FOR

Do not treat animals within one day of shearing. Do not apply orally in combination with organophosphates. Do not wet skin due to possible run-off. Do not apply to calves under 6 months old. They may show hypersensitivity to organophosphates.

FOR CONTROL OF FACE FLIES, FLEAS, LONG STAR TICKS ON DAIRY CATTLE: Mix 1 quart of CIODRIN VAPONA Insecticide (10% concentration). Spray to thoroughly 1 - 2 pints of spray per large animal. For pen or corral operations involving equipment, do not exceed a 0.5% Insecticide Emulsifiable Concentrate. Thoroughly cover the animal's body with 1 quart of spray per large animal and application as necessary to maintain improved face fly control. May be applied per 6 - 10 gallons of finished spray control, but not more often than once a week.

LOW VOLUME APPLICATION FOR CONTROL OF HORSES: Mix 1 quart of CIODRIN-VAPONA Insecticide (concentration). Using a calibrated sprayer, apply 5 - 10 teaspoons to the back and sides of the animal. This is achieved by the addition of 1 level teaspoon of water to each 100 ml of insecticide.

FOR CONTROL OF LICE AND WINTERSHAFTS AND HOGS: Mix 2 quarts of CIODRIN-VAPONA Insecticide (concentration) per 16 gallons of water (0.125% concentration). Using a calibrated sprayer, apply a second application to maintain control, but not more often than once a week.

BACKRUBBER APPLICATIONS: For horn flies and face flies, mix 1 quart of CIODRIN-VAPONA Insecticide (concentration) per 4 gallons of back rub oil or fuel oil in accordance with local regulations.

FOR CONTROL OF HOUSE FLIES, LIVESTOCK BUILDINGS: Mix 2 quarts of CIODRIN-VAPONA Insecticide (concentration) per 6 gallons of water (0.125% concentration). Spray outside barn walls (including dairy parlor) at least twice monthly using approximately 25% improved house fly and face fly solution at the rate of 1 pound per 6 gallons. Do not use in poultry houses.

FOR CONTROL OF CHORIOPTIC MITES: Mix 1 quart of CIODRIN-VAPONA Insecticide Emulsifiable Concentrate (concentration). Spray animals thoroughly.

NOTICE: Seller's guarantee shall be limited to the time period during which the buyer assumes any risk to persons or property. Seller does not accept the product on these conditions.

HP 2964

BEST DOG



CIODRIN VAPONA INSECTICIDE

Emulsifiable Concentrate

CONTAINS
CIODRIN and VAPONA Insecticides

ACTIVE INGREDIENTS	BY WEIGHT
Dimethyl phosphate of alpha-methyl-3-hydroxy-cis-erectone*	13.7%
2,2-dichlorovinyl dimethylphosphosphate	2.2%
Related Compounds	0.2%
Xylene	72.0%
TOTAL	100.0%

* CIODRIN® Insecticide
Equivalent 2.5% Vapona (DDVP) Insecticide

KEEP OUT OF REACH OF CHILDREN **WARNING**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED - CALL A PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY. Patient presents as usual toward if possible vomiting stomach contents and informed physician. Drink one or two glasses of water and induce vomiting by touching the back of the throat with a soft cloth or sponge or give anything by mouth to an adult or child except a physician.

IF INHALED. Remove victim to fresh air. Apply first aid respiration if indicated.

IF ON SKIN. Remove contaminant. Wash skin areas with soap and water.

IF IN EYES. Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

NOTE TO PHYSICIAN - ANTIDOTE STATEMENT
No specific antidote exists. Treatment is symptomatic and supportive and may be directed by a physician.

See side-back panel for additional precautionary statement

NET CONTENTS (LITERS)

EPA Reg No. 11842-9 EPA Est. No. 11842-A-2

Manufactured By
InterChem, Inc.
Alton, Iowa 51003

DIRECTIONS FOR LIVESTOCK PEST CONTROL

Do not treat animals with a history of sensitivity.
Do not apply directly to contact with the animal.
Do not wet sensitive areas of the animal.
Do not apply to a very underdeveloped calf or Brangus (Brangus cattle) until it is treated as they may show hypersensitivity to some insecticides.

FOR CONTROL OF FACE FLIES, HORN FLIES, HOUSE FLIES, STABLE FLIES, LICE AND LONG STAR TICKS ON DAIRY CATTLE, BEEF CATTLE, SHEEP, GOATS AND HOGS: Mix 2 quarts of CIODRIN-VAPONA Insecticide Emulsifiable Concentrate per 6 gallons of water (16% concentration). Spray thoroughly over all parts of the animal including the legs using 1-2 quarts of spray per large animal and up to 1 quart less for small animals. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

Improved face fly control may be obtained by applying the same solution at a rate of 1 pound per 6-10 gallons of finished spray. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

LOW VOLUME APPLICATION FOR CONTROL OF FACE FLIES: Mix 4 tablespoons (2 fluid oz.) of CIODRIN-VAPONA Insecticide Emulsifiable Concentrate per 1 pint of water (16% concentration). Using a hand held sprayer apply 5 teaspoons of spray to the face and a total of 5-10 teaspoons to the back and sides of the animal. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

FOR CONTROL OF LICE AND WINTER TICKS ON DAIRY CATTLE, BEEF CATTLE, SHEEP, GOATS AND HOGS: Mix 2 parts of CIODRIN-VAPONA Insecticide Emulsifiable Concentrate per 16 gallons of water (16% concentration). Spray animal thoroughly using up to 1 gallon of spray. Apply a second application 14 days later. Repeat applications as necessary to maintain control, but not more often than once every 7 days.

BACKRUBBER APPLICATIONS: For use in backrubbers or tail rubbers for the control of barn flies and face flies mix 1 quart of CIODRIN-VAPONA Insecticide Emulsifiable Concentrate per 4 quarts of backrubber or tail rubber. Dilute by adding fuel or fuel oil in a 1:1 ratio with the insecticide.

FOR CONTROL OF HOUSE FLIES, FACE FLIES, AND STABLE FLIES IN AND AROUND LIVESTOCK BUILDINGS: Mix 2 quarts of CIODRIN-VAPONA Insecticide Emulsifiable Concentrate per 8 gallons of water (16% concentration). Spray entire surfaces of and around farm walls (including dairy cattle houses), pens, and other structures containing animals at a rate of 1 quart per 8 gallons of water. Repeat above application as necessary to maintain control by applying 1 quart of spray per 8 gallons of water.

For use in poultry houses.

FOR CONTROL OF CHORIOPTIC MITES ON DAIRY AND BEEF CATTLE: Mix 2 parts of CIODRIN-VAPONA Insecticide Emulsifiable Concentrate per 16 gallons of water (16% concentration). Spray animal thoroughly. Retreat within 10-14 days for lasting control.

NOTICE: The user shall be bound by the terms of the label and subject the label to the distributor if any risk to persons or property arises out of misuse or improper handling and application, or just on these conditions.

HP 2964

BEST DOCUMENT AVAILABLE