



# 5% KORLAN INSECTICIDE GRANULES

CONTROLS Hog Lice on all ages of swine - newborn, too.

Also controls housefly Maggots (larvae), and chicken body lice on poultry.

<b>ACTIVE INGREDIENTS:</b>	
RONNEL (O,O-dimethyl O-(2,4,5-trichloro phenyl) phosphorothioate*)	5%
<b>INERT INGREDIENTS</b>	95%

<b>E.P.A. Registration No. 11849-13</b>
<b>E.P.A. Est. No. 11849-1A-2</b>
® Trademark of the Dow Chemical Company
<b>From technical grade; not suitable for internal use in animals.</b>

## CAUTION

**KEEP OUT OF THE REACH OF CHILDREN  
HARMFUL IF SWALLOWED**

Avoid contamination of feed, food and water.  
Avoid skin contact. Wash thoroughly after using.

**Note to Physician and Veterinarian:** Ronnel is a cholinesterase inhibitor. Atropine is an antidote. Treat symptomatically.

### LIVESTOCK REMEDY - NOT FOR HUMAN USE

This product is TOXIC to fish. Keep out of lakes, streams, or ponds.  
Apply this product only as specified on this label.

## USE PRECAUTIONS

Do not reuse empty container. Destroy it by perforating or crushing it. Bury it or discard it in a safe place away from water supplies.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

## DIRECTIONS

Distribute 5% Korlan Insecticide Granules with a mechanical spreader, scoop, or can.

**CONTROL OF HOG LICE:** Apply 5% Korlan Insecticide Granules evenly over the bedding for swine at the rate of 1/2 lb. per 100 ft. of area. No mixing with the bedding is necessary. One treatment will control the hog lice infestation when the animals use the bedding regularly. Re-treat if reinfestation occurs.

**CONTROL OF CHICKEN BODY LICE ON POULTRY:** Scatter 5% Korlan Insecticide Granules evenly over the litter at the rate of 1 lb. 100 ft. 5% Korlan Insect Granules may be used in broiler, layer, and replacement pullet houses. Re-treat if reinfestation occurs.

**CONTROL OF HOUSEFLY MAGGOTS (LARVAE): CAGED-LAYER POULTRY HOUSES:** Apply 5% Korlan Insecticide Granules at the rate of 1 lb. 100 ft. of area. Apply evenly over manure piles under layer cages, and over the floor area where flies breed. Re-treat as needed.



Manufactured by  
**SILAK COMPANY**  
Hempstead, Iowa 51238

**NET WEIGHT 10 LBS.**

RECEIVED