

### CAUTION

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention. Keep away from open flame. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water, or foodstuffs. If swallowed, immediately get medical attention. NOTE TO PHYSICIAN — Antropine is antidotal.

DESTROY USED CONTAINERS: Do not re-use for any purpose. Destroy containers by breaking and burying fragments in isolated area. Dispose of in a way as not to constitute a hazard, or contaminate potable water supplies.

### GENERAL DIRECTIONS

1.0% VAPONA Insecticide animal spray solution is a specifically prepared ready-to-use insecticide for use in conventional hand, power or automatic spray equipment, electric, fogging or misting equipment.

For the control of listed insects attacking animals described on this label. Apply in accordance with State and Federal recommendations. For young animals use 1/2 the adult dosage. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

Do not treat animals within one day of slaughter.

Do not apply dermally in combination with Trichlorofon.

EPA Reg. No. 11849-16  
EPA Est. No. 11849-1A-2

Net Contents



## 1% VAPONA® INSECTICIDE READY-TO-USE ANIMAL SPRAY SOLUTION

For the Control of Horn Flies, Stable Flies, Face Flies, Mosquitoes and House Flies by Direct Application on Dairy and Beef Cattle.

| ACTIVE INGREDIENTS                   | PERCENT BY WEIGHT |
|--------------------------------------|-------------------|
| 2,2-dichlorovinyl dimethyl phosphate | 0.93*             |
| Related Compounds                    | 0.07*             |
| Deodorized Kerosene                  | 49.50             |
| Mineral Oil                          | 49.50             |
|                                      | <hr/>             |
|                                      | TOTAL 100.00      |

\*Equivalent to 1% VAPONA Insecticide

### CAUTION!

KEEP OUT OF THE REACH OF CHILDREN

See side panels for additional cautions.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Manufactured by

**SILAK COMPANY, INC.**

Moscow, Iowa 51238

### USE AND DIRECTIONS FOR APPLICATION

Cattle, Dairy and Beef — Flies (Horn Fly, Stable Fly, Face Fly, House Fly) Mosquitoes

Apply 1 to 2 fluid ounces of 1.0% solution per animal as a mist spray daily with hand or automatic sprayer. Spray to thoroughly cover all parts of the animal, including the legs, but do not wet the skin. Do not apply in excess of 2 fluid ounces per day per animal. There is no pre-slaughter interval when used on beef animals.

ANIMAL BUILDINGS — Horse Stables, Dairy Barns, Auction & Show Barns (Flies, Gnats, Mosquitoes)

Apply 1.0% VAPONA Insecticide as a fog or mist. 1 pint of a 1% solution will treat 8000 cubic feet (40 X 20 X 10). In dairy barns, before milking, the fog or mist can be applied over and around each animal to kill flies. Before applying reduce air movement as much as possible by closing doors, windows and other openings. Do not use this solution for fogging or misting purposes in areas where animals have received a direct application within 8 hours. Do not contaminate water, feed or foodstuffs, milk or milking utensils.

MAY 27 1975