MAR 28 1975 Under the Frederic Andrew ELA May and

Poultry Houses: Scatter the dry bait or apply the liquid bait to areas where flies congregate, but keep out of reach of birds. Apply to manure, on window sills, outside of penned enclosures and on floor of feed storage room. Repeat as necessary Apply 1/4 lb. dry bait or 1 qt. liquid bait per 1,000 sq. ft.

Dog Kennels: Scatter the dry bait or apply the liquid bait outside dog runways, on window sills, ledges and other surfaces out of reach of animals. For caged units apply to dropping area. Repeat as necessary. Apply 1/4 lb. dry bait or 1 qt. liquid bait per 1,000 sq.ft.

Garbage Cans: Apply the dry bait around and inside garbage cans. Repeat as necessary.

CAUTION:

Harmful if swallowed, inhaled or absorbed through the skin.

Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Notice of Warranty

SILAK makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label

E.P.A. REG. NO. 11849-17 E.P.A.

E.P.A. Est. No. 11849-1A-2



SCATTER BAIT VAPONA INSECTICIDE

Use Wet or Dry

Active Ingredients	By
2.2-dichlorovinyl dimethyl phosphate.	•
Related Compounds	•
Inert Ingredients	
Total	
*Equivalent to 0.5%W of VAPONA Ins	

CAUTION: Keep out of reach of children. See side panel for additional cautions.

Do not reuse container. Destroy when empty.

Net contents: 2 POUNDS



MANUFACTURED BY: SILAK COMPANY, INC, BOX 173, HOSPERS, IOWA 51238

Use and Directions for Applications

Vapuna Insecticide Fly Bait is a specially prepared insecticidal bait for use in the control of flies and can be used wet or dry.

Liquid Bait: Dissolve 1 12 of bait in 1 gallon of water and apply as a coarse spray, sprinkle or spray.

Dry Bait: Scatter bait where flies congregate.

For reducing populations of Resistant and Non-resistant Houseflies in and around Out-door Living Areas, Picnic Grounds, Backyard Areas, Patios, Loading Docks, Outdoor Latrines, Parking Areas, Refuse Areas around Service Stations, Open-air Drive-ins, Ice Cream Stands, Garbage Collection and Disposal Areas: Scatter the dry bait or apply the liquid bait to areas where flies congregate. For open-air drive-ins, ice cream stands and restaurants, apply on apron between serving counter and parking area, paying particular attention to trash barrels and garbage cans where flies congregate. Apply 1/2 lb dry bait or 2 gt liquid bait per 1,000 sq. ft.

Food Processing Plants, Restaurants: Use only when the facility is not in operation and food is not exposed. Use only in bait boxes where bait in each box can be accounted for. Remove and account for all bait boxes prior to resuming food processing.

Dairy Barns, Milk Sheds, Stables, Livestock Barns, Loafing Sheds, Pig Pens and Outdoor Areas. Scatter the bait or apply the liquid bait to floors or ground areas where flies congregate. Ideal baiting sites are around doorways, feed storage rooms, alleyways and window sills. On freshly limed floors, place treatment on waste cloth or on paper. Repeat as necessary for practical fly control. Apply 1/4 lb dry bait or 1 qt. liquid bait per 1,000 sq. ft.

Caged Poultry and Turkeys on Wire: Scatter the dry bait or apply the liquid bait on droppings beneath the cages or perches Repeat at weekly intervals Apply 1/4 Ib dry bait or 1 gt liquid bait per 1,000 sg ft.

y Weight 0.46%* 0.04%* 99.50% 100.00%

secticide