

# MALATHION GRAIN COATING

A POWDERED INSECTICIDE FOR USE ON STORED GRAINS. Wheat, Corn, Oats, Rye, Barley, Grain Sorghum, Field and Garden Seeds.

PROTECTS AGAINST DAMAGE from Confused Flour Beetle, Rice Weevil, Granary Weevil, Saw toothed Grain Beetle, Flat Grain Beetle, Red Flour Beetle, Rusty Grain Beetle, Lesser Grain Borer, Indian Meal Moth.

<b>ACCEPTED</b>  <b>AUG 1 1974</b>  UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER- ED UNDER NO. 11849-18	<b>ACTIVE INGREDIENTS:</b> Malathion* . . . . . 2%
	<b>INERT INGREDIENTS:</b> . . . . . 98% *0, 0-dimethyl dithiophosphate of diethyl mercaptosuccinate

## CAUTION KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed. Avoid breathing of dust. Avoid contact with skin. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Apply this product only as specified on the label. Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.



## IMPORTANT

Use of this product as directed below will be within the tolerance of 8 parts per million for MALATHION. Treated grain is not subject to downgrading and can be milled into flour or feed using normal production methods.

## DIRECTIONS FOR USE

Apply MALATHION GRAIN COATING to the grain at the rate of 30 lbs. per 1000 bu. as the grain is being loaded or turned into final storage.

Field and garden seeds can be protected against grain pests

## HOW TO APPLY TO SHELLED CORN AND SMALL GRAINS

Apply MALATHION GRAIN COATING when grain is in the truck or wagon prior to binning. To determine the

feed and foodstuffs. Apply this product only as specified on the label. Do not reuse container. Destroy when empty.

See label

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

## IMPORTANT

Use of this product as directed hereon will not constitute a warranty, and it is not subject to downgrading and/or withdrawal.

## DIRECTIONS FOR USE

Apply MALATHION GRAIN COATING to grain at the rate of 30 lbs. per 1000 bu. as the grain is being loaded or turned into final storage.

Field and garden seeds can be protected against grain losses with a dosage of 30 lbs. of MALATHION GRAIN COATING per 1000 bu. of seed.

To protect stored grains from attack by Indian Meal Moth, apply MALATHION GRAIN COATING to the surface of clean or infested grain at the rate of 15 lbs. per 1000 sq. feet of grain surface area. Apply the dust evenly over the surface of the grain. Apply immediately after grain is loaded into storage and repeat if necessary.

## HOW TO USE SHELLE CORN AND SMALL GRAINS

Apply MALATHION GRAIN COATING to grain at the rate of 30 lbs. per 1000 bu. as the grain is being loaded or turned into final storage. Measure out the grain in a truck or wagon, and spread it evenly over the surface of the grain in the truck or wagon. Cut into the grain with a shovel or fork, so that the coating becomes mixed with grain as it is dumped from the truck or wagon into the hopper. Further mixing will occur when the grain is augered or blown into the bin. Mixing is completed as the coating and grain roll down the cone shaped pile in the bin.

CONTAINS  
**CYTHION**  
THE PREMIUM GRADE MALATHION  
EPA REG. NO. 11840-18



Manufactured by  
**SILAK COMPANY, INC.**  
Hospers, Iowa 51238

E.P.A. Reg. No. 11840-18  
E.P.A. Est. 11840-1A-2  
Net wt. 50 lbs.