



Master Mix®

RESEARCH APPROVED

Rabon*

50% WETTABLE POWDER

MASTER MIX RABON 50% WETTABLE POWDER is suitable for use in conventional power or low pressure knapsack sprayers. Occasional agitation is recommended to prevent undue settling of the suspension. Follow the USE DIRECTIONS for the proper percent solution needed for a specific insect and area of type of walls or surfaces. Refer to DILUTION TABLE for the quantity of insecticide needed to make the percent solution recommended.

		DILUTION TABLE		
		AMOUNT OF WATER		
		0.5% SOLUTION	1% SOLUTION	2% SOLUTION
5 Tablespoons		2 Gal	1 Gal	0.5 Gal
1/2 Lb		6 Gal	3 Gal	1.5 Gal
1 Lb		25 Gal	12.5 Gal	6.25 Gal
4 Lb		50 Gal	25 Gal	12.5 Gal
8 Lb		100 Gal	50 Gal	25 Gal

PRECAUTIONS IN USING

Harmful if swallowed. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of food and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or deposit of waste. Apply this product only as specified on this label. Do not reuse container. Destroy when empty.

NOTICE OF WARRANTY: CENTRAL SOYA makes no warranty or merchantability. fitness for any purpose, express or implied, concerning this product or its use, which extend beyond the statements on this label.

DIRECTIONS FOR USE					
TO CONTROL	PREVENTIVE AREAS	SCREEN SIZE	APPLICATION	REMARKS	
Fleas	Cavy barns poultry houses stable barns other animal buildings	20%	1 lb. 1 gal. 10 gal.	1 lb. 1 gal. 10 gal.	Apply wettable form to concrete floors surfaces. Form planted, weed or poison control blocks surfaces. Form mixture on galvanized steel metal surfaces.
Mosquitoes	Poultry droppings manure piles garbage piles order feed troughs	10%	1 gal.	1 gal.	Excessive control areas may require repeat applications.
Fleas	Camp grounds play yards, etc. areas around camp parks, other camping areas	20%	1 lb. 1 gal. 10 gal.	1 lb. 1 gal. 10 gal.	Apply treated areas thoroughly. Reapply treated areas and boundaries leading to treated areas.
Others	Swedes	0.5%	1 gal.	1 gal.	Apply to seedbeds, etc., to prevent infestation.
Others	Grasshoppers	0.5%	1 gal.	1 gal.	Apply to grasshopper infested areas.
Others	Locusts	0.5%	1 gal.	1 gal.	Apply to locust infested areas.
Others	Ants	0.5%	1 gal.	1 gal.	Apply to ant infested areas.
Others	Beetles	0.5%	1 gal.	1 gal.	Apply to beetle infested areas.
Others	Worms	0.5%	1 gal.	1 gal.	Apply to worm infested areas.
Others	Other insects	0.5%	1 gal.	1 gal.	Apply to other insect infested areas.

CENTRAL SOYA
MANUFACTURERS
OF AGRICULTURAL
CHEMICALS



CENTRAL SOYA
Feed Division

ACCEPTED

11/11/1976

Under the Federal Insecticide,
Fungicide and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 11886-6

BEST AVAILABLE COPY