

ACCREDITED

MAY 11 1977

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is registered as a disinfectant.

12015-14

10F2
Pm 31
12015-14

USE DIRECTIONS FOR SANITIZING

Sanitizing of Food Processing Equipment and other Hard Surfaces in Food Contact Locations.

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard surfaces. Wash and rinse all articles thoroughly, then apply a solution of 1 oz. **SPEEDEE 1000** in 4 gallons of water.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz./4 gallon dilution of **SPEEDEE 1000**.

Food contact surfaces must be rinsed with potable water before use.

At 1 oz./4 gallons, **SPEEDEE 1000** has a hard water tolerance of 400 ppm when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz. **SPEEDEE 1000** in 4 gallons of warm water.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

DANGER

Keep out of reach of children. **CORROSIVE** Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

12015HO1

SPEEDEE - 1000

Disinfectant-Sanitizer-Deodorizer with Organic Soil Tolerance

Disinfectant-Sanitizer-Deodorizer for Hospital, Institutional, Industrial, School, Dairy and Other Farm and Home Use.

AOAC Phenol Coefficients

<i>Staphylococcus aureus</i> , ATCC # 6539	100
<i>Salmonella typhosa</i> , ATCC #6539	73

Active Ingredients

Octyl decyl dimethyl ammonium chloride	5.0%
Didecyl dimethyl ammonium chloride	2.5%
Dioctyl dimethyl ammonium chloride	2.5%
Isopropyl alcohol	4.0%

Inert Ingredients

86.0%
100.00%

NET CONTENTS ONE GALLON

EPA Registration No. 12015-14

Do not reuse empty container

ALLIED LABORATORIES, INC.

875 Lake Rd.
Medina, Ohio 44256

DANGER:

**KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANELS FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.**

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

USE DIRECTIONS FOR DISINFECTION

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions.

For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces.

Add 3 1/2 oz. **SPEEDEE 1000** to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.

At this use-level, **SPEEDEE 1000** is effective against *Pseudomonas aeruginosa* when evaluated against *Pseudomonas aeruginosa* when evaluated by the AOAC Use-Dilution Test.

SPEEDEE 1000 is effective against *Staphylococcus aureus* and *Salmonella typhosa* even in the presence of organic soil and residual anionic detergents when used at 3 oz./5.5 gallons.

Disinfectant in Institutions, Industry, Schools and Homes.

For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces.

Add 3 oz. of **SPEEDEE 1000** to 5.5 gal. Apply to previously cleaned hard surface cloth.

At 3 oz./5.5 gallon use-level, **SPEEDEE 1000** is effective against *Staphylococcus aureus*, *Salmonella choleraesuis*.

Disinfection of Barber Tools
Barber tools (such as combs, brushes, scissors) can be disinfected by immersion in a 1/2-oz./gallon solution of **SPEEDEE 1000**.
Disinfection of Poultry Equipment, Animal Kennels.

Poultry brooders, watering troughs, equipment and other animal quarters (and kennel areas) can be disinfected by cleaning by applying a solution of 3 1/2 oz. **SPEEDEE 1000** to 4 gallons of water with a brush. Small utensils should be immersed in solution.

Prior to disinfection, all poultry, other their feeds must be removed from the area. This includes emptying all troughs, rack feeding and watering appliances. Remove and droppings from floors, walls and other occupied or traversed by poultry or other animals. After disinfection, ventilate buildings, other closed spaces. Do not house poultry animals, or employ equipment until feeds have been absorbed, set or dried.

All treated equipment that will come in contact with drinking water must be rinsed with water before reuse.

Rinse empty container thoroughly with water and discard it.

ED
1977
Insecticide,
Fungicide, Act,
Pesticide
2015-14

12015RH01

SPEEDEE - 1000

Disinfectant-Sanitizer-Deodorizer
with Organic Soil Tolerance

Disinfectant-Sanitizer-Deodorizer for Hospital, Institutional, Industrial, School, Dairy and Other Farm and Home Use.
DAC Phenol Coefficients
Staphylococcus aureus, ATCC # 6538 100
Salmonella typhosa, ATCC #6539 73

Active Ingredients
Sodium decyl dimethyl ammonium chloride 5.0%
Sodium lauryl dimethyl ammonium chloride 2.5%
Sodium dodecyl dimethyl ammonium chloride 2.5%
Isopropyl alcohol 4.0%

Inert Ingredients 86.0%
Water 100.00%

NET CONTENTS ONE GALLON

Registration No. 12015-14
Do not reuse empty container

ALLIED LABORATORIES, INC.
975 Lake Rd.
Medina, Ohio 44258

DANGER:

KEEP OUT OF REACH OF CHILDREN
READ SIDE PANELS FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

USE DIRECTIONS FOR DISINFECTION

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions.

For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces.

Add 3 1/2 oz. **SPEEDEE 1000** to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.

At this use-level, **SPEEDEE 1000** is effective against *Pseudomonas aeruginosa* when evaluated against *Pseudomonas aeruginosa* when evaluated by the AOAC Use-Dilution Test.

SPEEDEE 1000 is effective against *Staphylococcus aureus* and *Salmonella typhosa* even in the presence of organic soil and residual anionic detergents when used at 3 oz./5.5 gallons.

Disinfectant in Institutions, Industry, Schools and Homes

For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces.

Add 3 oz. of **SPEEDEE 1000** to 5.5 gallons of water. Apply to previously cleaned hard surface with mop or cloth.

At 3 oz./5.5 gallon use-level, **SPEEDEE 1000** is effective against *Staphylococcus aureus* and *Salmonella choleraesuis*.

Disinfection of Barber Tools

Barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2-oz./gallon solution of **SPEEDEE 1000**.

Disinfection of Poultry Equipment, Animal Quarters and Kennels.

Poultry brooders, watering founts, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 3 1/2 oz. **SPEEDEE 1000** to 4 gallons of water with a mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Rinse empty container thoroughly with water and discard it.