

DIRECTIONS FOR USE

DISINFECTION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply AQUAPHENATE with a cloth, mop, or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes, and other Health Care Institutions. For disinfecting floors, walls, counter tops, bathing areas, lavatories, bed frames, tables, chairs, garbage pails, and other hard, non-porous surfaces:

Add 3 1/2 oz. AQUAPHENATE to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth. At this use level, AQUAPHENATE is effective against Pseudomonas aeruginosa.

Disinfection in Institutions, Industry, and Schools. For disinfecting floors, walls, bed frames, counter tops, tables, chairs, garbage pails, bathroom fixtures, and other hard, non-porous surfaces:

Add 2 oz. AQUAPHENATE to 4 gallons of water. Apply to previously cleaned hard surface with mop or cloth. At 2 oz. / 4 gallon use-level, AQUAPHENATE is effective against Staphylococcus aureus, Salmonella choleraesuis, and Trichophyton mentagrophytes (the athlete's foot fungus).

Disinfection of Barber Tools. Pre-cleaned barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2 oz. / gallon solution of AQUAPHENATE.

Disinfection of Poultry Equipment, Animal Quarters, and Kennels. Poultry brooders, watering founts, feeding equipment, and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz. AQUAPHENATE in 4 gallons of water with a mop, cloth, or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all littering and droppings from floors, walls, and other surfaces occupied or traversed by poultry or other animals. After disinfection, ventilate buildings, coops, and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set, or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

SANITIZATION

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, counter tops, refrigerated storage and display equipment and other hard, non-porous surfaces. No potable water rinse is required. Wash and rinse all articles thoroughly, then apply a solution of 1 oz. Aquaphenate in 4 gallons of water. Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be reused for sanitizing applications.

Apply to sink tops, counter tops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush.

Directions for use in Eating Establishments:

- 1. Scrape and pre-wash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner (a commercial cleaner or detergent can be recommended).
3. Rinse with clean water.
4. Sanitize in a solution of 1/4 ounce to 1 gallon of water (150ppm). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drainboard to air dry.

Note: A clean potable rinse following sanitization is not permitted under Section HSS 196.13 of the Wisconsin Administrative Code.

At 1 oz. / 4 gallons, Aquaphenate fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinances 1978 Recommendations of the U.S. Public Health Services in water up to 1000 ppm of hardness calculated as CaCO3 when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz. Aquaphenate in 4 gallons of warm water. No potable water rinse is required.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

PRECAUTIONARY STATEMENTS -

Hazards to Humans and Domestic Animals

DANGER

KEEP OUT OF REACH OF CHILDREN. CORROSIVE. CAUSES EYE DAMAGE AND SKIN IRRITATION. DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING. PROTECT EYES AND SKIN WHEN HANDLING. HARMFUL IF SWALLOWED. AVOID CONTAMINATION OF FOOD.

AQUAPHENATE

Disinfectant-Sanitizer-Fungicide Deodorizer for Hospital, Institutional, Industrial, School, Dairy and Other Farm Use.

Table with 2 columns: Ingredient type and percentage. ACTIVE INGREDIENTS: Didecyl dimethyl ammonium chloride (7.5%), INERT INGREDIENTS (92.5%).

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No. 12017-3 EPA Establishment No. 12017-PA-1

Net Contents ONE GALLON



APTER INDUSTRIES

BOX 456, McKEESPORT, PA 15134-0456 U.S.A. (412) 872-9628 CALL TOLL FREE CONTINENTAL U.S.A. (800) 441-7146



ACCEPTED FOR CARE

NOV 21 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 12017-3

STORAGE AND DISPOSAL

- 1. Prohibitions: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
2. Storage Instructions: Store in original container. Keep from freezing.
3. Spill or Leak Procedures: Small spills may be mopped up, flushed away with water or absorbed on some absorbent material and incinerated. Large spills should be contained; the material then moved into containers and disposed of by approved methods for hazardous wastes.
4. Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the user according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.
5. Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
IF CONTAINER IS 1 GALLON OR LESS, DO NOT REUSE EMPTY BOTTLE. WRAP CONTAINER AND PUT IN TRASH.
6. General: Consult Federal, state or local disposal authorities for approval of alternative procedures such as limited burning.

COPYRIGHT © OCTOBER 1996 APTER INDUSTRIES, INC., McKEESPORT, PA. U.S.A.

BEST AVAILABLE COPY

2013

AQUAPHENATE

**Disinfectant-Sanitizer-Fungicide Deodorizer
for Hospital, Institutional, Industrial,
School, Dairy and Other Farm Use.**

ACTIVE INGREDIENTS

Didecyl dimethyl ammonium chloride 7.5%

INERT INGREDIENTS 92.5%

100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.


NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No. 12017-3

EPA Establishment No. 12017-PA-1

Net Contents **ONE GALLON**



APTER INDUSTRIES

BOX 456, McKEESPORT, PA 15134-0456 U.S.A.
(412) 672-9628

CALL TOLL FREE

CONTINENTAL U.S.A. (800) 441-7146

ACCEPTED
NOV 21 1990
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the registration of registered and...
12017-3
COLOR CARE SYSTEM

STORAGE AND DISPOSAL

1. Prohibitions: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

2. Storage Instructions: Store in original container. Keep from freezing.

3. Spill or Leak Procedures: Small spills may be mopped up, flushed away with water or absorbed on some absorbent material and incinerated. Large spills should be contained; the material then moved into containers and disposed of by approved methods for hazardous wastes.

4. Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

5. Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

IF CONTAINER IS 1 GALLON OR LESS, DO NOT REUSE EMPTY BOTTLE. WRAP CONTAINER AND PUT IN TRASH.

6. General: Consult Federal, state or local disposal authorities for approval of alternative procedures such as limited burning.

COPYRIGHT © OCTOBER 1996

APTER INDUSTRIES, INC., McKEESPORT, PA S.A.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

DISINFECTION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply AQUAPHENATE with a cloth, mop, or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes, and other Health Care Institutions.

For disinfecting floors, walls, counter tops, bathing areas, lavatories, bed frames, tables, chairs, garbage pails, and other hard, non-porous surfaces:

Add 3 1/2 oz. AQUAPHENATE to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth. At this use level, AQUAPHENATE is effective against *Pseudomonas aeruginosa*.

Disinfection in Institutions, Industry, and Schools.

For disinfecting floors, walls, bed frames, counter tops, tables, chairs, garbage pails, bathroom fixtures, and other hard, non-porous surfaces:

Add 2 oz. AQUAPHENATE to 4 gallons of water. Apply to previously cleaned hard surface with mop or cloth.

At 2 oz. / 4 gallon use-level, AQUAPHENATE is effective against *Staphylococcus aureus*, *Salmonella choleraesuis*, and *Trichophyton mentagrophytes* (the athlete's foot fungus).

Disinfection of Barber Tools.

Precleaned barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2 oz. / gallon solution of AQUAPHENATE.

Disinfection of Poultry Equipment, Animal Quarters, and Kennels.

Poultry brooders, watering founts, feeding equipment, and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz. AQUAPHENATE in 4 gallons of water with a mop, cloth, or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all littering and droppings from floors, walls, and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops, and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set, or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

SANITIZATION

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, counter tops, refrigerated storage and display equipment and other hard, non-porous surfaces. No potable water rinse is required. Wash and rinse all articles thoroughly, then apply a solution of 1 oz. Aquaphenate in 4 gallons of water, (150 ppm active). Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be reused for sanitizing applications.

Apply to sink tops, counter tops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush.

Directions for use in Eating Establishments:

1. Scrape and pre-wash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner (a commercial cleaner or detergent can be recommended).
3. Rinse with clean water.
4. Sanitize in a solution of 1/4 ounce to 1 gallon of water (150ppm). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drainboard to air dry.

Note: A clean potable rinse following sanitization is not permitted under Section HSS 196.13 of the Wisconsin Administrative Code.

At 1 oz. / 4 gallons, Aquaphenate fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinances 1978 Recommendations of the U.S. Public Health Services in water up to 1000 ppm of hardness calculated as CaCo3 when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz. Aquaphenate in 4 gallons of warm water. No potable water rinse is required.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

PRECAUTIONARY STATEMENTS — Hazards to Humans and Domestic Animals

DANGER

KEEP OUT OF REACH OF CHILDREN. CORROSIVE. CAUSES EYE DAMAGE AND SKIN IRRITATION. DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING. PROTECT EYES AND SKIN WHEN HANDLING. HARMFUL IF SWALLOWED. AVOID CONTAMINATION OF FOOD.

BEST AVAILABLE COPY