

Aug 22 1972
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR CONTROL OF PESTS
 REGISTERED UNDER
 SECTION 241

Cross-infection is of major housekeeping concern not only in hospitals, but in schools, institutions, and industry. SANISHINE is formulated for this problem area. It both cleans and disinfects effectively when used as directed. Its hard surface disinfecting action will reduce the hazard of cross-infection.

Two ounces of SANISHINE per gallon of water will deodorize bathrooms, garbage storage areas, and other areas where bacterial growth can cause mal odors.


WARNING

KEEP OUT OF REACH OF CHILDREN

Causes severe eye and skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink promptly a large quantity of water. Avoid alcohol. Call a physician immediately.



SANISHINE

**Cleaner — Disinfectant — Deodorizer — Fungicide
 for Hospital and Institutional Use**

AOAC Phenol Coefficients

Staph. aureus (ATCC No. 6538)	100
Salmonella typhosa (ATCC No. 6539)	50

Active Ingredients

Didecyl dimethyl ammonium chloride	4.5 %
Tetrasodium ethylenediamine tetraacetate	2.0 %
Sodium carbonate	1.0 %
Sodium metasilicate, anhydrous	0.5 %

Inert Ingredients

92.0 %
100.00 %

EPA Registration No. 12042-1

NET CONTENTS:

VERAX CHEMICAL COMPANY

MAITRY INDUSTRIAL PARK

P. O. Box 603

Bothell, Washington 98021

WARNING

KEEP OUT OF REACH OF CHILDREN.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

USE DIRECTIONS:

Apply SANISHINE to walls, floors, and surfaces such as tables, chairs, etc. with a cloth or mop. For heavy soiling preliminary cleaning may be necessary.

For Hospitals and Nursing Homes: Use one ounce per gallon of water.


At two ounces per gallon use is effective against Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella typhosa, and pathogenic E. coli. Performance against the first three has been confirmed by the AOAC. Fungicidal performance was T. interdigitale by the AOAC.

For Schools, Institutional and Public Buildings: Use one ounce per gallon of water.

At one ounce per gallon use delivers excellent cleaning and is effective against Staphylococcus aureus, Salmonella choleraesuis, Escherichia coli, and phyton interdigitale, the same AOAC tests used above. Performance for hospitals was confirmed.

Rinse empty container with water.

Aug 7 1972
 UNDER THE FEDERAL INSECTICIDE
 ACT AND THE FEDERAL FUNGICIDE ACT
 FOR CONTROL OF PESTS
 REGISTERED BY THE



SANISHINE

**Cleaner — Disinfectant — Deodorizer — Fungicide
 for Hospital and Institutional Use**

AOAC Phenol Coefficients

Staph. aureus (ATCC No. 6538)	100
Salmonella typhosa (ATCC No. 6539)	50

Active Ingredients

Didecyl dimethyl ammonium chloride	4.5 %
Tetrasodium ethylenediamine tetraacetate	2.0 %
Sodium carbonate	1.0 %
Sodium metasilicate, anhydrous	0.5 %

Inert Ingredients

92.0 %
100.00 %

EPA Registration No. 12042-1

NET CONTENTS:

VERAX CHEMICAL COMPANY

**MAITRY INDUSTRIAL PARK
 P. O. Box 603
 Bothell, Washington 98021**

WARNING

**KEEP OUT OF REACH OF CHILDREN.
 SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY
 STATEMENTS**

housekeeping concern
 in schools, institutions,
 is formulated for this
 ns and disinfects effec-
 ed. Its hard surface dis-
 ce the hazard of cross-

IE per gallon of water
 , garbage storage areas,
 cterial growth can cause

**IING
 CH OF CHILDREN**

cin irritation. Do not get
 othing. Harmful if swal-
 ion of food.

diately flush eyes or skin
 at least 15 minutes. For
 Remove and wash all
 fore reuse. If swallowed,
 quantity of water. Avoid
 immediately.

USE DIRECTIONS:

Apply SANISHINE to walls, floors and other hard surfaces such as tables, chairs and bed frames with a cloth or mop. For heavily soiled areas, a preliminary cleaning may be required.

For Hospitals and Nursing Homes: Add two ounces per gallon of water.

At two ounces per gallon use-level, SANISHINE is effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella choleraesuis*, *Escherichia coli*, and pathogenic fungi. Germicidal performance against the first four organisms has been confirmed by the AOAC Use-Dilution Test. Fungicidal performance was determined against *T. interdigitale* by the AOAC Fungicidal Test.

For Schools, Institutional and Industrial Use: Add one ounce per gallon of water.

At one ounce per gallon use-level, SANISHINE delivers excellent cleaning and is germicidal. It is effective against *Staphylococcus aureus*, *Salmonella choleraesuis*, *Escherichia coli*, and *Trichophyton interdigitale*, the athlete's foot fungus. The same AOAC tests used above to confirm suitable performance for hospitals were used.

Rinse empty container with water before discarding.