

DANGER

Keep out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard.

WARRANTY

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials. Statements concerning the use of this product are not to be construed as recommending the infringement of any patent, and no liability for infringement arising out of any such use is assumed.

EPA Est. 12042-WA-1



DISINFECTANY - DEODORIZER - SANITIZER

EPA Reg. No. 12042-4

Active Ingredients	
n-Alkyl (60% C _{14,} 30% C _{16,} 5 - C _{2,} 5% C ₁₈)	
dimethyl benzyl ammonium chlorides	5%
n-Alkyl (68% C _{12,} 32% C ₁₄) dimethyl	
ethylbenzyl ammonium chlorides	. 5%
Inert Ingredients	90%
	100%
This product is a concentrate and should be diluted before usi	ng.

DANGER

(See other precautionary and first aid statements on left side of panel.)

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: ONE GALLON (3.77 L.)

USE DIRECTIONS:

Cold Disinfection

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1 ½ ounces per gallon of water for 10 minutes.

Food Processing Equipment

For sanitization of previously cleaned food processing equipment and food utensils dilute 1 ounce per 4 gallons of water to provide 200 ppm of active quaternaries. At this level, **No Potable Water Rinse** is required.

Dairies

To sanitize dairy equipment such as tanks, lines, pails, and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce per 4 gallons of water to provide 200 ppm of active quaternaries. At this level, **No Potable Water Rinse** is required. Consult local Health Board for additional information.

This product when used on environmental, inanimate, hard surfaces at 3½ ounces per 5 gallons of water is effective against Influenza A₂, Herpes Simplex, Adenovirus type 2, and Vaccinia Viruses.



Maltby Industrial Park Maltby, Washington 98290