

POISON

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

PRO - BOLL

M.P.-4

Contains 4 pounds Methyl Parathion per gallon

ACTIVE INGREDIENTS:

O, O-Dimethyl O-p-Nitrophenyl Phosphorothioate	45.48%
Xylene	49.52%

INERT INGREDIENTS: 5.00%

*Technical Methyl Parathion (80%)

____ GALLONS NET

DANGER!

KEEP OUT OF REACH OF CHILDREN

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME



See antidote statement and other precautions on back/side panel


NOTICE OF WARRANTY

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Manufactured by

PRO-BOLL CHEMICAL COMPANY, INC.
DUNN, LOUISIANA

ESTABLISHMENT No. 12062-A-1




WARNING SYMPTOMS APPEAR: Administer two 1/100 grain atropine tablets immediately. NEVER ADMINISTER UNLESS WARNING SYMPTOMS APPEAR (See WARNING SYMPTOMS below) Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps

TREATMENT: Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine as the specific therapeutic antagonist of choice against parasympathetic stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until the time for cholinesterase regeneration has been allowed as determined by blood test.



EPA REG. NO. 12062-5

DIRECTIONS FOR USE

When preparing a spray solution of PRO-BOLL M.P.-4, fill spray tank approximately $\frac{1}{2}$ full with water. Then with agitation (mechanical or overflow) in operation, add measured amount of PRO-BOLL M.P.-4. Then finish adding desired amount of water. Apply solution uniformly for most efficient results.

COTTON: Red Spider Mites, Boll Weevils, Leafworms, Aphids, use $\frac{1}{2}$ pint per acre in sufficient water to cover. Repeat at 3-day intervals. Do not hand pick or harvest within 7 days of application. Workers entering fields within 24 hours after application should wear protective clothing.

GRAINS (Wheat, Oats, Barley): Greenbugs (Aphids), use $\frac{1}{2}$ to 1 pint per acre in sufficient water to cover. Repeat as necessary. Do not apply after heads begin to form.

Do not feed the treated plants or trash to dairy animals or animals being fattened for slaughter.

PRECAUTIONS IN USING

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Decontaminate and destroy empty containers to prevent re-use. Decontaminate by washing thoroughly with a strong solution of alkali and detergent and rinse with water.

DANGER

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO METHYL PARATHION POISONING.

Do not swallow, get in eyes, on skin, or on clothing. Do not breathe dust or vapor. Wear clean natural rubber gloves; clean, waterproof protective clothing, and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type approved by the U. S. Bureau of Mines for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If material gets in eyes, immediately flush with water for at least ten minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before re-use. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrate solution of alkali and detergent and rinse with water.

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR METHYL PARATHION POISONING. Consult your physician about obtaining an adequate supply of 1/100 grain atropine tablets for emergency use. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from vicinity of the Methyl Parathion, remove all contaminated clothing, and wash skin with soap and water for at least ten minutes.

