

DIRECTIONS FOR USE

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Do not remain in treated areas. Ventilate the area after treatment is completed.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control *roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worms, grain mites, and cadelles:* Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control *carpet beetles:* Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets, and infested areas of shelving. Repeat treatment as needed.

To control *bedbugs:* Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings, and floors. Repeat treatment as needed.

To control *fleas and brown dog ticks:* Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, windows and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control *clothes moths:* Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages, directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.



SHERWOOD



GENERAL PURPOSE INSECTICIDE

A Concentrated PYRENONE® Micro Emulsion

– Clear

– Stable

– Leaves No Oily or Objectionable Residue

Effective, Fast Acting Insecticide.

For Use in Homes, Institutions, Restaurants.

Food Processing Plants, Dairies, Livestock and Outdoors for Mosquito Control.

Apply with Conventional Mechanical or Compressed Air Equipment.

ACTIVE INGREDIENTS:

Pyrethrins 0.1%

*Piperonyl Butoxide, technical 1.0%

Petroleum distillate 0.4%

INERT INGREDIENTS: 98.5%

*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and related compounds. .2%

Pyrenone is a registered trademark of FMC Corporation

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

Net Contents

Gallon


EPA Reg. No. 12123-3

EPA Est. No. 12123-NJ-1

MANUFACTURED BY

SHERWOOD CHEMICALS LTD

WESTVILLE, N.J. 08093



SPACE SPRAYING: shut off ventilator machine adjusted for rapid control of *skipper flies, weevils, moths, Mediterranean moths, and tobacco* upward angle distribute the entire area, at least 5 feet of space.

FOGGING: To kill the above named insects, 1 gallon to 20,000 square feet. Contact insects die.

OUTDOOR APPLICATION: as a fine mist to control insects. For best results, do not exceed 5 miles.

Treat shrubbery and trees. Shrubbery may rest. Shrubbery pools, marshy areas are treated. Application water is prohibited.

Apply at a rate of 1000 square feet).

Using truck-mount sprayer such as Rotomist traveling at 5 m.p.h. calibrate equipment.

Harmful if swallowed and foodstuffs.

Remove pets, birds and spraying.

ENVIRONMENTAL PRECAUTIONS

This product is toxic to fish and aquatic life. Do not apply to streams and ponds. Do not apply to areas where drift is likely to occur. Do not apply to areas where drift is likely to contaminate water. Do not apply to areas where drift is likely to contaminate water. Do not apply to areas where drift is likely to contaminate water.

PRECAUTIONS: Before storage or handling, be strict about cleanliness. Avoid