



50% SEVIN® CONCENTRATE DUST

CONTAINS SEVIN® CARBARYL INSECTICIDE

FOR THE CONTROL OF INSECTS ON COTTON

CAUTION:

KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENT :

Carbaryl (1-naphthyl N-methylcarbamate)..... 50%

INERT INGREDIENTS:

50%
50%
100%

*Trade Mark of Union Carbide Corporation.

DO NOT RE-USE EMPTY CONTAINER. Destroy it by Burying with Waste or Burning. Stay Away From Smoke or Fumes.

DIRECTIONS FOR USE ON COTTON

DIRECTIONS FOR USE IN THE EASTERN UNITED STATES ONLY

For early-season thrips, flea beetles, cotton fleahoppers, cotton leafworms, and striped blister beetles: Apply 2 lbs. of 50% Sevin Concentrate Dust per acre on appearance of insects.

For mid- and late-season bollworms, fall armyworms, cotton leaf perforators, leaf rollers, leafhoppers, and tarnished plant bugs. Also for light-to-moderate infestations of lygus bugs in Western irrigated cotton: Apply 4 lbs. of 50% Sevin Concentrate Dust per acre, depending on the size of the cotton and insect population level. Applications should be made every 5 to 7 days as long as control is necessary.

For pink bollworms, boll weevils, stink bugs, and salt marsh caterpillars: Apply 5 lbs. of 50% Sevin Concentrate Dust per acre on a 5 to 7 day schedule depending on the size of the cotton and degree of infestation.

For grasshoppers: Apply 2 to 3 lbs. of 50% Sevin Concentrate Dust per acre on appearance of insects and their feeding damage. The highest dosage should be used when grasshoppers are mature or material is applied on partly defoliated plants.

Aphid populations will be suppressed by repeated applications of this insecticide.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

NOTE

1. Calibrate ground equipment or aerial dust apparatus accurately. After properly adjusting equipment to deliver the required low volume, maintain constant speed throughout application of Concentrate SEVIN Dust to insure even coverage. Strictly avoid excessive application.

To effectively apply Concentrate SEVIN Dust by aircraft use standard venturi type equipment with spreader vanes or Swathmaster type spreaders. Fly at normal height (4 to 6 feet above the top of the plants is suggested) and maintain regular swath width (usually 30 to 35 feet is indicated). For application with ground equipment use tractor mounted power clusters with a metering device that feeds the Concentrate SEVIN Dust through one nozzle per row with deflector) or similarly designed electronic dusters.

For optimum results apply Concentrate SEVIN Dust only when weather conditions are favorable. Wind and rising air currents may cause undesirable drift. Strictly avoid excessive dust drift away from crop being treated.

SEVIN does not control spider mites. Where infestations are encountered, use a recommended miticide.

5. SEVIN may be applied after bolls open. Allow 7 days between last application and grazing treated fields or harvesting crop residue for use as feed or bedding for dairy or meat animals.

6. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

7. Special concentrate dusting equipment which can accurately apply a dosage as low as 1 pound per acre must be used to apply this Concentrate Dust.

Screen Size Specifications

Through 100 mesh (dry screening)	minimum 99.5%
Through 200 mesh (dry screening)	minimum 70.0%
Through 325 mesh (dry screening)	maximum 70.0%

Average particle size, microscopic analysis, between 30 and 60 microns.

CAUTIONS

HARMFUL IF INHALED OR SWALLOWED. AVOID BREATHING OF DUST. DO NOT TAKE INTERNALLY. SKIN CONTACT MAY BE HARMFUL. Avoid contact. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Manufactured By

FARM CHEMICALS, INC.

Aberdeen, North Carolina

Raeford, North Carolina

Net Contents

Lbs.

EPA Reg. No. 12130-2

ACCEPTED
11/4/73
UNION CARBIDE INSECTICIDE
FUNCTIONAL INSECTICIDE ACT
FOR ENVIRONMENTAL REGISTER
EPA UNDER 12130-2
NO ATTACHED COMMENTS

12130-2