

DANGER

Poisonous by skin contact, inhalation or swallowing! Rapidly absorbed through skin! Do not get in eyes, on skin or clothing. In case of skin contact, immediately remove all contaminated clothing and flush skin or eyes with plenty of water for at least 15 minutes. For eyes, get medical attention. Do not breathe dust or spray mist. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

Wear clean, weather-resistant clothing. Do not eat, drink or use tobacco while using this product. For more information, see Material Safety Data Sheet.

ANTIDOTES: If swallowed - give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a physician immediately. If on skin - wash immediately with soap and water.

DANGER: Do not use indoors or in other poorly ventilated places. Do not apply or allow to drift where dairy animals or animals being finished for slaughter may be grazing. Unprotected humans and beneficial animals should be kept out of treated areas for at least an hour after treatment. When workers must enter fields within 5 days of a spray they should be protected. The Federal Food and Drug Administration has set official tolerances for pesticide residue. Crops having residue in excess of these tolerances should not be marketed without removal of such excess.

POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish and birds and other wildlife. Do not feed or treat animals with this product. Keep out of streams of water. Do not apply when weather conditions may cause drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of water. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Protect from temperature below 32° F.

Rinse equipment and containers and dispose of wastes and spillage by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes or by burning (keep out of smoke).

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

NOT FOR USE OR STORAGE IN OR AROUND THE HOUSE.

DO NOT STORE NEAR OR CONTAMINATE FEED OR FEED STUFF.

NET CONTENTS

GALLONS

ACCEPTED

Feb 23, 1973

UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND RODENTICIDE
FOR ECONOMIC POISON RESISTANCE
EPA UNDER NO. 12130-5

Farm Chem

12130-5

ENDRIN 1.6-E

(Contains 1.6 pounds Endrin per gallon)



POISON



DANGER: KEEP OUT OF THE REACH OF CHILDREN

See Side Panel For Additional Warnings
And Antidote Statements

ACTIVE INGREDIENTS:

Endrin (Hexachloroepoxyoctahydro-Endo,

Endo-dimethanonaphthalene)

19.48%

Xylene

77.84%

INERT INGREDIENTS:

2.68%

100.00%

manufactured by

FARM CHEMICALS, INC.

ABERDEEN, NORTH CAROLINA

RAEFORD, NORTH CAROLINA

E.P.A. REG. NO. 12130-5

DIRECTIONS FOR USE

Put the required amount of this material through screen into partially filled spray tank with agitator running. Keep agitator running while mixing and spraying. If mixture does not mix readily but tends to separate as an oily layer, do not use as injury to plants may result.

NOTE: For airplane application, use the indicated dosage in 3 to 10 gallons of water per acre, depending on the stage of growth and amount of foliage to be covered. For conventional ground equipment, use 10 to 50 gals. of water per acre.

WHEAT, OATS, RYE AND BARLEY: Army Cutworms and Pale Western Cutworms - Use 1 pt. (0.2 lbs. of actual Endrin) per acre in sufficient water for good coverage. Single application. Do not pasture animals in treated fields. Do not apply within 45 days of harvest. Do not feed threshings to livestock.

COTTON: Thrips, Cotton Fleahoppers, Rapid Plant Bug, Tarnished Plant Bug - use 0.2 lbs. actual Endrin per acre (1 pint Endrin 1.6E). Apply in sufficient water for full coverage.

BOLL WEEVIL, COTTON BOLLWORM: For early season control use 0.2 lbs. actual Endrin per acre. For mid-season and late season control use 0.2 to 0.5 lb. actual Endrin per acre. (To obtain a rate of 0.5 lb. Endrin per acre, use 2 1/2 pints Endrin 1.6E per acre.)

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions and under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not readily foreseeable to seller, and buyer assumes the risk of any such use.

RECEIVED

Feb 23, 1973

FOR INFORMATION OF THE
OFFICE OF THE ATTORNEY GENERAL
AND THE DEPARTMENT OF
CORRECTIONS
ATTACHING DOCUMENTS.



100-30-5

ENDRIN

100-30-5

APPEAR TO ITS BYE AND DAILY

100-30-5

100-30-5

100-30-5