



SEVIN* 5 DUST

ACTIVE INGREDIENT - Carbaryl (1-naphthyl N-methylcarbamate)
INERT INGREDIENTS

5% by wt.
95% by wt.
100%

U.S. Pat. Nos. 2,903,478 3,009,855

Trade Mark of Union Carbide Corporation for the active ingredient carbaryl insecticide

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

DO NOT RE-USE EMPTY CONTAINER. Destroy it by Burying with Waste or Burning. Stay Away From Smoke or Fumes.

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

Preharvest and Grazing Use Limitations

No post treatment time limitations on applications to alfalfa, beans, blueberries, carrots, okra, clovers, peppers, corn, sorghum, or further to corn, eggplants, forage, grasses, garden peas, grapes, melons, peanuts, pears, potatoes, pumpkins, sugarcans, grown for forage, soybeans, sugar, sweet corn, and tomatoes.

NOTE

Recommended dosage refer to units of 50% dust per acre. The surface must be kept to obtain full coverage. Apply when insects or other damage is present. Repeat at 7 to 10 day intervals or as necessary.

Some phytotoxicity may occur on shade foliage in the presence of rain or high humidity of several days duration. Follow with irrigation.

For information on grazing use of SEVIN on pastures and forages, consult your local Agricultural Extension Office.

SMALL FRUIT INSECT CONTROL

BLACKBERRIES, BOYSENBERRIES, RASPBERRIES, LOGANBERRIES AND RASPBERRIES - Use 40 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

BEANS - Use 20 to 40 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

BLUEBERRIES - Use 40 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

CRANBERRIES - Use 20 to 40 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

GRAPES - Use 20 to 40 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

STRAWBERRIES - Use 20 to 40 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

VEGETABLE INSECT CONTROL

SPINACH - Use 20 to 40 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

BEANS - Green beans, Lima beans, cran beans, soybeans and black-eyed peas. Use 20 to 40 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

Use 20 to 40 pounds for armyworms, cabbage caterpillars, corn earworm, tarnished plant bug, and tobacco hornworm. Use 40 pounds for optimum control of horn earworm after lettuce heads begin to form. Use 40 pounds for optimum control of tobacco hornworm. Where cabbage looper is the principal pest, use an alternative pesticide. For control of small cabbage loopers after edible portions begin to form, use SEVIN in a 5 to 7 day schedule.

Allow 3 days between application and harvest of head lettuce, garden beets, broods, horseradish, parsnips, rutabagas, salsify (roots), and turnips (roots).

Allow 14 days between last application and harvest of Chinese cabbage, heads, dandelion, endive, escarole, garden beets, tops, kale, leaf lettuce, mustard greens, radishes, and turnips (tops), chard and turnips (tops).

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI, Use 10 to 20 pounds for flea beetle and harlequin bug. **Allow 3 days** between application and harvest.

Use 20 to 40 pounds for armyworms, cabbage caterpillars and corn earworm. Where cabbage looper is the principal pest, use an alternative pesticide that is specifically recommended. For control of small cabbage loopers after edible portions begin to form, use SEVIN in a 5 to 7 day schedule.

TOBACCO INSECT CONTROL

In plant beds: For tobacco flea beetle, apply 1 to 2 pounds per 100 square yards when insects appear. Excessive application plant injury may result. Repeat as needed at 10 to 14 day intervals and just before transplanting.

Injury may result when used on small plants or with application of heavy dosages on tender foliage during or when humidity remains high for several days following application of SEVIN. When dusting with a shaker or can, be sure not to overdose. Rotary dusters are recommended.

In fields: Broadcast applications for armyworms and tobacco flea beetles, apply 15 to 25 pounds per acre depending on size of the plants. Good coverage is essential. Use the highest dosage only on large plants and repeat as needed. Heavier applications may cause plant injury.

For late applications, allow three days before pruning or cutting. Treated fields may be entered immediately after dust has settled.

FORAGE AND FIELD CROP INSECT CONTROL

ALFALFA, CLOVERS, COWPEAS, PEANUTS AND SOYBEANS - Use 10 to 20 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

Use 20 pounds for alfalfa caterpillar, bean leaf beetle, cucumber beetle, green stink bug, and soybean leafhopper. Use 20 to 30 pounds for armyworms, corn earworm, stink bugs and white worms.

To avoid possible injury on tender foliage, do not apply when foliage is wet or when dew is present. Do not apply during the next two days.

FORAGE GRASSES AND PASTURE - Use 10 to 20 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

Use 10 to 20 pounds for armyworms, corn earworm, stink bugs and white worms. Use 20 to 30 pounds for armyworms, corn earworm, stink bugs and white worms.

SORGHUMS (including MILO AND GRAIN SORGHUM) - Use 20 to 40 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

Use 20 to 40 pounds for armyworms, corn earworm, stink bugs and white worms. Use 20 to 40 pounds for armyworms, corn earworm, stink bugs and white worms.

GRASSHOPPERS

Apply 10 to 20 pounds per acre. Apply 2 weeks before harvest. Repeat at 10 to 14 day intervals or as necessary.

Use 10 to 20 pounds for armyworms, corn earworm, stink bugs and white worms. Use 10 to 20 pounds for armyworms, corn earworm, stink bugs and white worms.

CAUTION

POISONOUS IF SWALLOWED OR INHALED

Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly after use. Avoid breathing dust.

Do not use in or near buildings, barns, or other structures. Do not use in or near water.

Note for Physician: Carbaryl is a carbamate. It is rapidly metabolized and excreted in the urine. It is not known whether it is a carcinogen.

AVOID CONTAMINATION OF FEED, FEEDING TROUGH, AND WATERING RECEPTABLES

NOTICE

12130-13
ACCEPTED
JAN 10 1973
FEDERAL BUREAU OF INVESTIGATION
U.S. DEPARTMENT OF JUSTICE

ras, berries. Allow 7 days between application to poultry and game birds or poultry and game and premises treatments and bird slaughter. Allow 7 days between last application and slaughter of beef cattle, horses, sheep and pigs.

BLUEBERRIES: Use 10 to 40 pounds per acre to control midget cherry and cranberry fruitworms and Japanese beetle. Apply 3 weeks before harvest and repeat 10 days later if necessary.
CRANBERRIES: Use 30 to 50 pounds per acre to control fruitworms, Japanese beetle and leafhoppers. Apply in late bloom and as needed thereafter. Allow 7 days between application and harvest.
GRAPES: Use 10 to 40 pounds per acre to control grape leafhopper and grape leaf skeletonizer. Apply just before first bloom and as needed thereafter. An extra application is needed for leafhoppers. Use 40 pounds for grape berry moth, Japanese beetle, grape beetles, grape leafhopper.
STRAWBERRIES: Use 10 to 40 pounds per acre to control strawberry root weevil, strawberry weevil and strawberry webworm. Allow 1 day between application and harvest.

VEGETABLE INSECT CONTROL

ASPARAGUS: Use 10 to 40 pounds per acre to control asparagus beetle, asparagus moth, asparagus fly, asparagus sawfly, asparagus webworm, asparagus root weevil, asparagus root borer, asparagus root miner, asparagus root girdler, asparagus root nematode, asparagus root rot, asparagus root blight, asparagus root wilt, asparagus root canker, asparagus root gall, asparagus root cyst, asparagus root nematode, asparagus root rot, asparagus root blight, asparagus root wilt, asparagus root canker, asparagus root gall, asparagus root cyst.
BEANS: Green beans, Lima beans, Snap beans, Snap peas and black-eyed peas. Use 10 to 40 pounds per acre to control bean beetle, bean fly, bean weevil, bean root weevil, bean root borer, bean root miner, bean root girdler, bean root nematode, bean root rot, bean root blight, bean root wilt, bean root canker, bean root gall, bean root cyst.
CORN: Use 10 to 40 pounds per acre to control corn earworm, corn rootworm, corn root borer, corn root miner, corn root girdler, corn root nematode, corn root rot, corn root blight, corn root wilt, corn root canker, corn root gall, corn root cyst.
PUMPKIN AND SQUASH: Use 10 to 40 pounds per acre to control pumpkin and squash beetle, pumpkin and squash fly, pumpkin and squash weevil, pumpkin and squash root weevil, pumpkin and squash root borer, pumpkin and squash root miner, pumpkin and squash root girdler, pumpkin and squash root nematode, pumpkin and squash root rot, pumpkin and squash root blight, pumpkin and squash root wilt, pumpkin and squash root canker, pumpkin and squash root gall, pumpkin and squash root cyst.
EGGPLANT AND PEPPER: Use 10 to 40 pounds per acre to control eggplant and pepper beetle, eggplant and pepper fly, eggplant and pepper weevil, eggplant and pepper root weevil, eggplant and pepper root borer, eggplant and pepper root miner, eggplant and pepper root girdler, eggplant and pepper root nematode, eggplant and pepper root rot, eggplant and pepper root blight, eggplant and pepper root wilt, eggplant and pepper root canker, eggplant and pepper root gall, eggplant and pepper root cyst.
CHINESE CABBAGE, CABBAGES, BROCCOLI, CAULIFLOWER, KALE, COLESLAW, GARDEN BEET, HORSERADISH: Use 10 to 40 pounds per acre to control Chinese cabbage, cabbages, broccoli, cauliflower, kale, coleslaw, garden beet, horseradish. Control insects and diseases.

Net Contents

Lbs.

GRASSHOPPERS

Apply 10 to 30 pounds as often as necessary to control grasshoppers on the above crops. A 10 to 20 pound rate is suggested for orchards, pastures, or sparse vegetation on waste and fringe lands, ditches and borders. Apply to 4-foot designations of the row when grasshoppers are mature or material is applied to the products of the row.

CAUTION

HARMFUL IF SWALLOWED OR INHALED.

Avoid contact with skin. Avoid contact with eyes. Skin Contact May Be Harmful.

Avoid contact with clothing. Avoid contact with food, feed, feeding troughs and watering receptacles. Avoid contact with children and animals.

Note for Physician: In case of poisoning, call physician immediately. Do not induce vomiting. Avoid contamination of food, feed, feeding troughs and watering receptacles.

NOTICE

Seller warrants that the product conforms to its official description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the seller, and any remedy shall be limited to any one of the following:

Manufactured By

FARM CHEMICALS, INC.

Liberty, N. C.

Branches in various states