



**CAN KILL YOU**

# 6-3 ETHYL-METHYL

An Emulsifiable Liquid

### ACTIVE INGREDIENTS

O, O-dimethyl O p nitrophenyl phosphorothioate 95

Parathion O O diethyl O p nitrophenyl phosphorothioate 10

INERT INGREDIENTS 26

### RECOMMENDATIONS VEGETABLES

DO NOT APPLY WITHIN 10 DAYS OF HARVESTING WITH

DO NOT APPLY WITHIN 10 DAYS OF HARVESTING WITH  
CORN, Seed Corn, Etc.

DO NOT APPLY WITHIN 10 DAYS OF HARVESTING WITH  
Beans and Turnips. Do not apply to any other vegetable  
crops.

CABBAGES

CUCUMBERS

PEAS

PEPPERS

DO NOT APPLY WITHIN 10 DAYS OF HARVESTING WITH  
CUCUMBER, KALE, MELON, AND

SPINACH

BEETROOT

DO NOT APPLY WITHIN 10 DAYS OF HARVESTING WITH  
CUCUMBER, KALE, MELON, AND

COTTON

PEAS

PEPPERS

PEPPERS

PEPPERS

PEPPERS

PEPPERS

PEPPERS

PEPPERS

PEPPERS

PEPPERS

PEPPERS

PEPPERS

011730

011730

APR 23



Form 226 (Rev. 10-12-65)

A.C. 100-100  
10/15/73  
12/10/73

phosphoramide Parathion (O, O diethyl O p nitropheno- l phosphorothioate INERT INGREDIENTS	28.95 57.89 13.26
--	-------------------------



# FARM CHEMICALS, INC.

ABERDEEN, WAKE COUNTY, NORTH CAROLINA

E.P.A. REG NO 12130-25

**COTTON:**  
Insecticide for use on cotton in the bollworm stage. Apply to the undersides of leaves. Do not apply more than 100 lbs per acre in a season.

**SUGAR BEETS:**  
Apply to the undersides of leaves. Do not apply more than 100 lbs per acre in a season.

**SORGHUM:**  
Apply to the undersides of leaves. Do not apply more than 100 lbs per acre in a season.

**TObACCO:**  
Apply to the undersides of leaves. Do not apply more than 100 lbs per acre in a season.

**IF YOU ARE USING THIS PRODUCT ON A CROP OTHER THAN COTTON, SUGAR BEETS, SORGHUM, OR TOBACCO, CHECK THE DIRECTIONS ON THE PRODUCT LABEL.**

NOTICE