

FOSTRACIDE

General Disinfectant Deodorizer
Fomexol, Fostacidin
Sanitizer with Soap - Sanitizing
Hospital, Institutional, and Industrial Use

Active Ingredients

Ethyl Dimethylammonium Chloride	4.0
Tetrasodium ethylene diamine tetraacetate	1.0
Sodium carbonate	1.0
Sodium metasilicate trihydrate	11.5

Inert Ingredients

12.1
100.0

**ONLY FOR SALE TO, USE AND STORAGE
BY SERVICE PERSONNEL
KEEP OUT OF REACH OF CHILDREN**

WARNING

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of water, egg whites, gelatin solution. Avoid alcohol.

Call a physician immediately.

**SEE LEFT PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS.**

EPA Registration No. 12179-5
EPA Establishment No. 12179 MI 1
Net Contents

FOSTER CHEMICALS INCORPORATED

DETROIT, MICHIGAN 48236

Directions for Use

GENERAL CLASSIFICATION

In violation of Federal Law to use this product in a manner inconsistent with its labeling.

FOSTRACIDE is a disinfectant, deodorizer, sanitizer with soap, and industrial use.

FOSTRACIDE is a disinfectant, deodorizer, sanitizer with soap, and industrial use.

Directions for Use: For general disinfection, dilute with water.

FOSTRACIDE is a disinfectant, deodorizer, sanitizer with soap, and industrial use.

Sanitizing: For sanitizing, dilute with water.

In hospitals and nursing homes, add 2.5 oz. per gallon of water.

FOSTRACIDE has been evaluated in the laboratory and is effective against the following organisms:

Staphylococcus aureus, Staphylococcus epidermidis, Bacillus subtilis, Pseudomonas aeruginosa, Escherichia coli, Salmonella typhimurium, Shigella flexneri, and Shigella sonnei.

AGAC Fungicide Test: **FOSTRACIDE** has been evaluated in the laboratory and is effective against the following organisms:

Aspergillus niger, Aspergillus fumigatus, Penicillium chrysogenum, and Penicillium notatum.

For use in medical institutions and industrial uses: **FOSTRACIDE** is a disinfectant, deodorizer, sanitizer with soap, and industrial use.

Sanitizing Non-Food Contact Surfaces: such as walls, floors, tables, and furniture.

FOSTRACIDE is a disinfectant, deodorizer, sanitizer with soap, and industrial use.

FOSTRACIDE is a disinfectant, deodorizer, sanitizer with soap, and industrial use.

FOSTRACIDE is a disinfectant, deodorizer, sanitizer with soap, and industrial use.

Two uses of FOSTRACIDE: For general disinfection, dilute with water.

Precautionary Statements

Hazards to Humans and domestic animals:

WARNING

Keep out of reach of children. May cause severe eye irritation or eye damage. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

STORAGE AND DISPOSAL

THIS PRODUCT MUST BE KEPT UNDER LOCKED STORAGE SUFFICIENT TO MAKE IT INACCESSIBLE TO CHILDREN OR PERSONS UNFAMILIAR WITH ITS PROPER USE.

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

OPEN DUMPING IS PROHIBITED.

DO NOT REUSE EMPTY CONTAINERS.

PESTICIDE DISPOSAL

PESTICIDE SPRAY MIXTURES: BEWARE THAT THIS PRODUCT IS NOT TO BE USED ON OR IN THE AREA WHERE IT IS USED. SHOULD BE DISPOSED OF IN A MANNER APPROVED FOR PESTICIDES OR BURIED IN A PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL

TRIPLE RINSE FOR COMPLETE REMOVAL OF RESIDUE. REGENERATE OR REUSE AS APPROVED. DO NOT REUSE FOR OTHER PURPOSES.

GENERAL

CONSULT FEDERAL, STATE, OR LOCAL HEALTH AUTHORITIES FOR APPROVED METHODS OF DISPOSAL OF SUCH AS EMPTIES, FOR A REUSE.

Cross contamination is of major housekeeping concern not only in hospitals, but in schools, institutions, and industry. **FOSTRACIDE** is formulated for this problem area. It both cleans and disinfects effectively and is virucidal when used as directed. Its hard surface disinfecting action will reduce the hazard of cross contamination.

FOSTRACIDE is an extremely effective one step sanitizer for non-food contact surfaces. It cleans and sanitizes in waters up to 850 ppm hardness. This hard water tolerance is in excess of most potable waters and will insure complete sanitizing action when used as directed.

Two ounces of **FOSTRACIDE** per gallon of water will deodorize surfaces in bathrooms, garbage storage areas, and other areas where bacterial growth can cause mal odors. Prevents odor caused by mold and mildew.

Precautionary Statements

Hazards to Humans and domestic animals

WARNING

Keep out of reach of children. May cause severe eye irritation or eye damage. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

STORAGE AND DISPOSAL

THIS PRODUCT MUST BE KEPT UNDER LOCKED STORAGE SUFFICIENT TO MAKE IT INACCESSIBLE TO CHILDREN OR PERSONS UNFAMILIAR WITH ITS PROPER USE.

—DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL

—OPEN DUMPING IS PROHIBITED

—DO NOT REUSE EMPTY CONTAINER

PESTICIDE DISPOSAL

PESTICIDE, SPRAY MIXTURE OR RINSATE THAT CAN NOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL

TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN A INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS OR BURY IN A SAFE PLACE.

GENERAL

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES SUCH AS LIMITED OPEN BURNING.

FOSTRACIDE

Cleaner-Disinfectant-Deodorizer-
Fungicide-Virucide*
Sanitizer with Organic Soil Tolerance for
Hospital, Institutional and Industrial Use

Active Ingredients

Didecyl dimethyl ammonium chloride	4.5%
Tetrasodium ethylenediamine tetraacetate	1.9%
Sodium carbonate	1.0%
Sodium metasilicate, anhydrous	0.5%

Inert Ingredients

92.1%
100.0%

"ONLY FOR SALE TO, USE AND STORAGE
BY SERVICE PERSONNEL"
KEEP OUT OF REACH OF CHILDREN

WARNING

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of water, egg whites, gelatin solution. Avoid alcohol.

Call a physician immediately.

SEE LEFT PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No. 12179-5
EPA Establishment No. 12179-MI-1
Net Contents

LIA Reg. No. 12179-5

Directions for Use

GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FOSTRACIDE is a proven one step cleaner-disinfectant sanitizer for use in the presence of moderate amounts of organic soil.

Apply **FOSTRACIDE** to walls, floors and other hard (non-porous) surfaces such as tables, chairs, countertops, sinks and fixtures with a cloth, mop, or mechanical spray device so as to thoroughly wet surfaces. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or when use of the solution becomes visibly dirty.

Disinfection — To disinfect hard, non-porous hospital surfaces, add 2 1/2 oz. per gallon of water. Treated surfaces must remain wet for 10 minutes. At this use level, **FOSTRACIDE** is also virucidal and fungicidal. At 1/2 oz. per gallon, **FOSTRACIDE** will disinfect hard surfaces in homes, schools, industry, and non-medical institutions.

Sanitizing — To sanitize porous or non-porous, non-food contact surfaces, add 1 oz. per 2 1/2 gallons of water. Treated surfaces must remain wet for 30 seconds.

For hospitals and nursing homes, add 2.5 oz. per gallon of water. The minimum activity of **FOSTRACIDE** has been evaluated in the presence of 5% serum and found to be effective against the following organisms by the AOAC Use-Dilution test:

- Pseudomonas aeruginosa
- Staphylococcus aureus
- Salmonella enteritidis
- Escherichia coli
- Streptococcus pyogenes
- Proteus mirabilis
- Klebsiella pneumoniae
- Enterobacter aerogenes
- Salmonella schottmuelleri
- Streptococcus faecalis
- Shigella dysenteriae
- Brevibacterium ammoniagenes

AOAC Fungicidal Test

FOSTRACIDE is an effective fungicide against Trichophyton mentagrophytes (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas, and telephones, etc.

Virucidal Performance

At 2 oz. per gallon use level, **FOSTRACIDE** was evaluated in the presence of 5% serum and found to be effective against the following viruses: herpes simplex, a member of the virus family that causes the common cold; a representative of the poliovirus; Adenovirus Type 5, a causative virus in respiratory diseases; and influenza A, as represented by the strains commonly known as Hong Kong Flu and the London Flu Virus, or influenza A virus, (H3N2).

For schools, non-medical institutional and industrial uses: At 1/2 oz. per gallon of water, **FOSTRACIDE** has been evaluated for germicidal effectiveness. It is effective against Staphylococcus aureus and Bacillus subtilis. The AOAC tests conducted at the University of Michigan hospitals were carried out in the presence of 5% serum.

Sanitizing Non-Food Contact Surfaces (such as walls, floors, tables, etc.) At 1/2 oz. per gallon use level, **FOSTRACIDE** is an effective sanitizer in waters of up to 850 ppm hardness against Staphylococcus aureus and Klebsiella pneumoniae in the presence of 5% serum on hard, non-porous surfaces.

FOSTER CHEMICALS INCORPORATED

DETROIT, MICHIGAN 48238

TEL (313) 867-5444