

**FOSTRACIDE EX** is a germicide with an organic soil tolerance. Its disinfecting action has been verified in the AOAC Use Dilution Test against *Staphylococcus aureus*, *Salmonella choleraesuis*, and *Pseudomonas aeruginosa* in the presence of 5% blood serum as a source of organic matter. This means that one step disinfection and cleaning can be achieved when used as directed.

Cross-contamination is of major housekeeping concern not only in hospitals, but in schools, institutions, and industry. **FOSTRACIDE EX** is formulated for this problem area. It cleans, disinfects, and is virucidal when used as directed. Its hard surface disinfecting action will reduce the hazard of cross contamination.

**FOSTRACIDE EX** is virucidal against Herpes Simplex (a member of the Herpes viruses that cause chicken pox, severe brain infections, and infectious mononucleosis), Vaccinia (representative of the pox viruses) and Influenza A<sub>1</sub> (the Hong Kong flu virus) on inanimate environmental surfaces.

#### Precautionary Statements

**Hazards to Humans and domestic animals**

### **DANGER**

Keep out of reach of children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed or absorbed through the skin. Avoid contamination of food.

#### STORAGE AND DISPOSAL

- DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL
- OPEN DUMPING IS PROHIBITED
- DO NOT REUSE EMPTY CONTAINER

#### PESTICIDE DISPOSAL

PESTICIDE THAT CANNOT BE USED, OR CHEMICALLY REPROCESSSED SHOULD BE DISPOSED OF IN A LAND FILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

#### CONTAINER DISPOSAL

DISPOSE OF IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS. BURY IN A SAFE PLACE OR RETURN TO DRUM RECONDITIONER.

#### GENERAL

SEE LEFT PANEL FOR FEDERAL, STATE, OR LOCAL DISPOSAL AND USES FOR APPROVED ALTERNATIVE PROCESSES. LIMITED OPEN BURNING.

**FOSTRACIDE EX**  
**Heavy Duty Germicide**  
**with Organic Soil Tolerance**  
Cleaner-Disinfectant-Deodorizer-Fungicide-  
Virucide-Sanitizer  
for Hospital and Institutional Use

<b>Active Ingredients</b>	
Octyl Decyl Dimethyl Ammonium Chloride	4.50%
Dioctyl Dimethyl Ammonium Chloride	2.25%
Didecyl Dimethyl Ammonium Chloride	2.25%
Tetrasodium Ethylenediamine Tetraacetate	2.28%
<b>Inert Ingredients</b>	88.72%
	100.00%

**KEEP OUT OF REACH OF CHILDREN**  
**"ONLY FOR SALE TO, USE AND STORAGE**  
**BY SERVICE PERSONNEL"**  
**DANGER**

**Statement of Practical Treatment**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short-acting barbiturate drug.

SEE LEFT PANEL FOR  
**ADDITIONAL PRECAUTIONARY STATEMENTS**

EPA Registration No.	12179-7
EPA Establishment No.	12179-MI-1
<b>Net Contents</b>	

#### Directions for Use

#### GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**FOSTRACIDE EX** is a proven one step cleaner-disinfectant sanitizer fungicide virucide in the presence of moderate amounts of organic soil when used as directed.

Apply **FOSTRACIDE EX** to walls, floors and other hard inanimate non porous surfaces such as tables, chairs, countertops, sinks and bedframes with a cloth, mop or mechanical spray device so as to thoroughly wet surfaces. For heavily soiled areas a preliminary cleaning is required. Prepare a fresh solution daily or when use solution becomes visibly dirty.

**Disinfection** — for routine disinfection of hard non porous surfaces, add 1 oz. per gallon of water. Treated surfaces must remain wet for 10 minutes. At this use level, **FOSTRACIDE EX** is also virucidal and fungicidal. When encountering organic soil in hospitals, nursing homes, schools, institutional and industrial uses, add 2 oz. per gallon of water.

**Sanitizing** — To sanitize porous or non porous, non food contact surfaces, add 2 1/2 oz. per gallon of water. Treated surfaces must remain wet for 60 seconds.

For hospitals, nursing homes, schools, institutions and industry, add 1 oz. per gallon of water. The broad spectrum activity of **FOSTRACIDE EX** has been evaluated in the presence of 5% serum and found to be effective against the following organisms by the AOAC Use Dilution test:

- Pseudomonas aeruginosa*
- Staphylococcus aureus*
- Salmonella choleraesuis*

#### AOAC Fungicidal Test

**FOSTRACIDE EX** is an effective fungicide against *Trichophyton mentagrophytes* (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas, exercise facilities, etc.

#### Virucidal Performance

At 1 oz. per gallon use level, **FOSTRACIDE EX** was evaluated in the presence of 10% serum and found to be effective against the following viruses: Herpes Simplex (a member of the virus family that causes infectious chicken pox), Vaccinia (representative of the pox viruses) and Influenza A<sub>1</sub> (represented by the strain commonly called the Hong Kong flu) on inanimate environmental surfaces.

**Sanitizing Non-Food Contact Surfaces** (such as walls, floors, tables, etc.): At 2 1/2 oz. per gallon use level, **FOSTRACIDE EX** is an effective sanitizer in waters of up to 400 ppm hardness against *Staphylococcus aureus* and *Klebsiella pneumoniae* in the presence of 5% serum on hard porous and non porous environmental surfaces.

**FOSTER CHEMICALS INCORPORATED**

DETROIT MICHIGAN 48238

TEL (313) 867-5444