# **Banish Microbiocide**

#### **ACTIVE INGREDIENT:**

This product contains 0.87 lb. of active ingredient per gallon and weighs 8.68 lb. per gallon.

## KEEP OUT OF REACH OF CHILDREN WARNING

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARRING: Causes eye demage and skin irritation. Harmful if swallowed. Do not get in eyes, on skin, or on clothing. Avoid contamination of food. Weer gougles or face shield and rubbar gloves when handling. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT: In case of skin contact, wash with plenty of soap and water Remove contaminated clothing and wash before reuse, if product gets in the eyes, flush immediately with copious amounts of clean, cool water for at least 15 minutes. Get medical attention immediately. If product is swallowed, call a physician immediately. If petient is conscious, induce vomiting by stroking or tickling the patient's throat or far back on patient's tongue. Emetics such as 2 teespoonsful (10 mL) of specae syrup or 1 teespoonful (5 mL) of dry mustard in warm water to form a paste or even soap in warm water can be used. Repeat until vomit fluid is clear. Then have patient drink plenty of milk, gelatin solution, beaten egg whites, flour and water, or other nonoily demul cent. Never induce vomiting or give anything by mouth to an unconscious person.

Note to physician Probable mucosal damage may contraindicate gastric lavage.

ENVIRONMENTAL HAZARDS: This pasticide is toxic to fish. Do not discharge treated effluent into lakes streams, ponds, or public waters unless in accordance with a NPDES permit. For guidance, contact your Regional Office of the Environmental Protection Agency.

Manufactured by:
Allstates Chemical Corporation
P. O. Box 619
Katy, TX 77449

EPA REG. NO. 12264-24

LPA EST. NO.

NET CONTENTS .....

### DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Banish Microbiocide is used in industrial and or commercial recirculating cooling tower systems and industrial air washing systems to control microbiological slime. Prior to the use of Banish Microbiocide in industrial and/or commercial recirculating cooling tower systems, systems should be cleaned to remove algal growth, microbiological slime, and other deposits. Then make an initial slug addition of 12.0 to 17.9 ff. oz. of Banish Microbiocide per 1000 gal, of water to provide 100 to 150 ppm of Banish Microbiocide, based on fill weight of water in the system. Repeat initial dosage until control is evident. Make subsequent slug addition of 6.0 to 17.9 ff. oz. of Banish Microbiocide per 1000 gal, of water (50 to 150 ppm Banish Microbiocide) every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleadoff and the severity of the microbiological problem. Slug additions should be made in the sump of recirculating cooling tower systems.

Banish Microbiocide is used in industrial air-washing systems which maintain effective mist-eliminating components. Prior to its use, systems should be cleaned to remove bacterial slime and other deposits. An initial slug dose of 28.1 to 35.9 fl. oz. of Banish Microbiocide per 1000 gal. of water is recommended. Repeat initial dosage until control is evident. Subsequent slug additions of 21.0 to 35.9 fl. oz. of Banish Microbiocide per 1000 gallons of water should be employed every 1 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and severity of the bacterial problem. Slug additions may be made to the sump or the water collection trays of the airwash system.

#### STORAGE AND DISPOSAL

m

口

U

PROHIBITIONS: Divinit containing at ewater flood, or feed by storage or disposal. Open dumping is prohibited PESTICIDE DISPOSAL. Pestigide: spray mixture, or it safe that cannot be used or chemically reprocessed should be disposed of in a landfle approved for pesticides, or buried in a safe place away from water supplies. CONTAINER DISPOSAL.

METAL Triple rinse for equivalents and other for recycling, re-onditioning, or disposation an approved landfalor body in a safe place. Containers over 30 gains is should be resealed before offering for reconditioning. PEASTIC Containers under 30 gains in ust not be reused but should be triple rinsed and disposed of in an incorrector or andfill approved for pest. I decenture ers or buried in a safe place. Containers over 30 gains are to be reserved as a feed of the containers over 30 gains are to be reserved as a feed for reconditioning or triple in section equivalents and offered for reconditional procedure.

GENERAL Considered on State of Colored Spassolar to State Colored Statement of smooth area.

- 1