[12367-4]

DIRECTIONS

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

The above described application will kill accessible, exposed stages of flies, including fruit flies, mosquitoes, silver fish, small flying moths, gnats, crickets, roaches, spiders, scorpions, wasps, hornets, clover mites, cheese mites, cheese skipper, and these common stored product pests; granary weevils, rice weevils, confused flour beetles, saw toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, Angoumois grain moths, Mediterannean flour moths, Indian meal moths, Tobacco moths, meal worms, and grain mites.

IN MUSHROOM PRODUCTION AND PROCESSING: To control adult Phorids, Sciards, and Cecidomylids in growing houses, apply at the rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned off. In packing and canning plants, use at the rate of 1 ounce to 2000 cubic feet of space.

CAUTION: Harmful if swallowed. Avoid contamination of feed and food-stuffs.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label. Do not reuse empty container. Destroy by perforating or cruching and burying in a safe place.

FOOD PLANT FOGGING INSECTICIDE

EPA Reg. No. 12367-4

A Concentrated PYRENONE® Insecticide

Designed for use in Edible Product Areas of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs, and in Other Food Processing Plants.

Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist.

ACTIVE INGREDIENTS:	100%
Pyrethrins	0.5%
Piperonyl Butoxide, Technical	4.0%
Petroleum Distillate	95.5%

*Equivalent to 3.2% (butylcarbityl) (6-propylpiperonyl) ether and to 0.8% of related compounds.

CAUTION KEEP OUT OF REACH OF CHILDREN

See rear panel for additional precautions

Net Contents: 1 Gallon

PYRENONE is a registered trademark of FMC Corporation

MANUFACTURED IN U.S.A. BY



SANITARY SUPPLIES/CLEANING MATERIALS/MAINTENANCE EQUIPMENT

- aza £ank 6.6 + 928 Lysle Blvd + McKeesport Pa 1513∂

123614