

ACCEPTED  
 132455-1  
 JUN 7 1974  
 UNDER FEDERAL REGULATION AS A PESTICIDE  
 FUNGICIDE AND ANTHERIZONE ACT  
 FOR ECONOMIC POISON REGISTRATION  
 ED UNDER NO. ----- SUBJECT  
 TO ATTACHED COMMENTS.

# "RAZE"<sup>TM</sup>

## RAT AND MOUSE BAIT

READY TO USE BAIT STATION FOR RATS AND MICE

**ACTIVE INGREDIENTS:**

Warfarin, (3-(Alpha-Acetylbenzyl)-4-Hydroxycoumarin), . . . . . 0.025%  
 N1-(2-Quinoxaliny) Sulfanilamide (Sulfaquinoxaline) . . . . . 0.025%

**INERT INGREDIENTS** . . . . . 99.950%  
 TOTAL 100.000%

**CAUTION**  
 KEEP OUT OF REACH OF CHILDREN  
 See additional cautions on back panel

EPA EST. NO. 12455-WI-1

NET WEIGHT 3 OZ.

EPA REG. NO. 12455-1

**DIRECTIONS**

**RATS:** Place at least 2 "RAZE" bait stations next to burrows, in runways, along walls, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 15 days and continue baiting until all signs of feeding have stopped.

Placing bait stations in concealed locations where rats and mice can feed unobserved is desirable.

**MICE:** Place "RAZE" bait stations where mice or signs of mice have been seen. Stations of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days.

Replace stale, damp or moldy bait with fresh bait stations.

This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for Warfarin.

**CAUTION**

Baits should be placed in areas not accessible to children, pets, wildlife and domestic animals or placed in tamperproof bait boxes.

If swallowed by humans, pets, domestic animals or wildlife, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. In case of ingestion, call a physician.

Do not reuse empty container, destroy it by burying with waste or burning. Stay away from smoke or fumes.

Manufactured by **BELL LABORATORIES, INC.** Box 5133 Madison, Wis. 53705