

... available

12455 -15

06/21/1977

DIRECTIONS

~~PEL~~ M... ..

Place baits in locations where rats and mice feed, water, and/or travel. Bait placements along walls, in corners and in concealed locations where rats and mice can feed unobserved are desirable. Contaminated or spoiled baits should be replaced promptly.

**NORWAY AND ROOF RATS:** Place 4 to 16 oz. of 'Final' bait in stations next to burrows, in runways, along walls, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 10 days and continue baiting until all signs of feeding have stopped.

**HOUSE MICE:** Place 1/4 to 1/2 oz. of 'Final' bait in stations 8 to 12 feet apart where mice or signs of mice have been seen. Assure an uninterrupted supply of bait for a period of not less than 15 days and continue baiting until all signs of feeding have stopped.

This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K.

# Final

## RAT and MOUSE BAIT



# Pelleted

**READY TO USE BAIT FOR RATS AND MICE**

### CAUTION

**KEEP OUT OF REACH OF CHILDREN**

See additional cautions on back panel

#### ACTIVE INGREDIENTS:

- Warfarin, 3-(alpha-Acetylbenzyl)-4-hydroxycoumarin ----- 0.025%
- N<sup>1</sup>-(2-Quinoxaliny) sulfanilamide (Sulfiquinoxaline) ----- 0.025%
- INERT INGREDIENTS** ----- 99.950%

TOTAL 100.000%

NET WEIGHT 1 lb. (454 g.)

**ACCEPTED**

JUN 21 1977

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 12455-15

### CAUTION

Pelleted baits must be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamperproof boxes. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Keep out of lakes, streams, or ponds. Do not reuse empty container, destroy it by burying with waste or burning. Stay away from smoke or fumes. If swallowed by humans, pets, domestic animals or wildlife, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of 4-hydroxycoumarin. In case of ingestion, call a physician.



Manufactured by  
**BELL LABORATORIES, INC.**  
 Box 2153, Madison, Wis. 53705, U.S.A.

EPA EST. NO. 12455-15