

RECEIVED
22 JAN 1980

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Keep away from humans, domestic animals, and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K₁ intramuscularly or orally, as indicated in baby-droxy-coumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD

Keep out of lakes, streams or ponds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway rats, roof rats, and house mice in and around homes, industrial and agricultural buildings, and similar manmade structures. Treated baits must be placed in locations not accessible to children, pets, domestic animals, or wildlife, or in tamper resistant bait boxes. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats, mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

P.C.Q.[®]

**RAT AND MOUSE BAIT
KILLS NORWAY RATS, ROOF
RATS, AND HOUSE MICE**

ACTIVE INGREDIENTS	2-Diphenylacetyl-L-proline	99.95%
INERT INGREDIENTS		0.05%
TOTAL		100.000%

CAUTION

KEEP OUT OF REACH OF CHILDREN
(See left panel for additional precautionary statements)

NET CONTENTS: 25 lbs.

Manufactured by:

B Bell Laboratories, Inc.
2001 Lenoir Avenue
Kenilworth, N.J. 07033

EPA EST. NO. 12455-19-1 EPA REG. NO. 12455-19

APPLICATION DIRECTIONS

RATS: Apply 3 to 16 oz of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 15 days.

MICE: Apply 1 to 2 oz (1 to 2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 oz) may be needed at points of high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS/MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store only in original container in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY