DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

 Store product not in use in a location out of reach of children and pets.

- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
- Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice in an `around homes, industrial and agricultural buildings, and similar m. -made structures.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS:

Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, patween floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

Ditrac

RAT AND MOUSE PAIT

KILLS NORWAY RATS, ROOF RATS, AND HOUSE MICE

KEEP OUT OF REACH OF CHILDREN CAUTION

(See right panel for additional precautionary statements)

Net Contents: 25 lbs.

Mig. by:



EPA EST. NO. 12455-WI-1

EPA REG. NO. 12455-19

APPLICATION DIRECTIONS

RATS: Apply 3 to 16 oz. of bait per placement. Maintain an ininterrupted supply of fresh bait for at Jeast 40 days.

MICE: Apply 1/4 to 1/2 oz. (1 to 2 level tablespoors) of bain at 8 to 12-foot intervals. Larger placements (uū to 2 ozil may be neerled at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permane it bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. S. TORAGE: Store only in original container, in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting form the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS CAUTION

Keep away from humans, domestic animals, and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleading.

NOTE TO PHYSICIAN: It ingested, administer Vitamin K, intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD

Keep out of lakes, streams or ponds.

ACCEPTED NOV 79 1992

Undor the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Res. No. 12455-19

BEST AVAILABLE COP"