

Reg. # 12455-67

PM-14

197

**ACCEPTED**  
 NOV 9 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 12455-67

# Ditraac®

## Mouse Bait Station

### Kills Mice

**ACTIVE INGREDIENT:**

Diphacinone;  
 2-(Diphenylacetyl)-1,3-indandione ..... 0.005%  
 INERT INGREDIENTS:.....99.995%  
 TOTAL 100.000%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

NOTE TO PHYSICIAN: If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

SEE LABEL ON OUTER PACKAGE FOR ADDITIONAL PRECAUTIONARY INFORMATION.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See outer package label for complete DIRECTIONS FOR USE including USE RESTRICTIONS and BAITING DIRECTIONS.


EPA Reg. No. 12455-67

EPA Est. No. 12455-WI-1

Manufactured by:



**Bell Laboratories, Inc.**  
 Madison, WI 53704 U.S.A.



of 3

Reusable

Prebaited

# Ditrac<sup>®</sup>

## Mouse Bait Station

### KILLS MICE


Active Ingredient:	
Diphacinone;	
(2-Diphenylacetyl-1,3-indandione) .....	0.005%
Inert Ingredients.....	99.995%
Total	100.000%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

(See back panel for additional precautionary statements)

**ACCEPTED**  
**NOV 9 1992**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 12455-67



#### DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### USE RESTRICTIONS:

THIS IS NOT A TAMPER-PROOF OR TAMPER-RESISTANT BAIT STATION. DO NOT USE THIS PRODUCT IN AREAS ACCESSIBLE TO CHILDREN, PETS, DOMESTIC ANIMALS, AND NONTARGET WILDLIFE. For control of house mice in and around homes, industrial, commercial, agricultural, and public buildings. Do not place in areas where there is possibility of contaminating food or surfaces that come in direct contact with food.

#### SELECTION OF TREATMENT AREAS:

Determine areas where mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, beside burrows, or in corners and concealed places.

#### APPLICATION DIRECTIONS:

**HOUSE MICE:** Place bait station at intervals of 8 to 10 feet along walls or in corners where mice live or travel. When bait is consumed, it may be replaced with one DITRAC All-Weather BLOX. (EPA Reg. No. 12455-80) Do not use any other bait block in this station.

**TO REFILL:** (See illustrations below) 1. Fully insert 2-pronged key in holes on top of DITRAC Mouse Bait Station. 2. Bend key away from station. 3. Gently lift up. 4. Place one DITRAC All-Weather BLOX in the triangular area next to the hinge. 5. Close lid until it snaps securely. Repeat steps for easy rebaiting.



**NET CONTENTS: 1 oz. (28 gms)**

EPA EST. NO. 12455-WI-1

EPA REG. NO. 12455-67

Manufactured by:



Bell Laboratories, Inc.  
Madison, WI 53704 • U.S.A.

To prevent reinfestation, eliminate food, water and harborage as much as possible. If reinfestation does occur, repeat treatment.

#### ENVIRONMENTAL HAZARDS

Keep out of lakes, streams and ponds.

#### STORAGE AND DISPOSAL:

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store only in original container in a cool, dry place; inaccessible to children and pets.

**DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

#### PRECAUTIONARY STATEMENTS:

Hazards to humans and domestic animals.

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician at once.

**NOTE TO PHYSICIANS:** If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.