

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will frequently find and consume the bait. Generally these are along walls, by ground openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

RATS: Apply 2 to 10 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.
HOUSE MICE: Place 1 Place Pac at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.
RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and remove bait regularly. To prevent reinfestation, limit sources of rodent food water and harborage as much as possible. Reinfestation does occur. Repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container in a dry area inaccessible to children and pets.
DISPOSAL: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

ACCEPTED
MAR 30 1992
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 12455-75

Contrac
RAT and MOUSE BAIT

READY TO USE PLACE PAC
KILLS RATS AND MICE

KILLS WARFARIN RESISTANT NORWAY RATS
 Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins

ACTIVE INGREDIENT Bromadiolone: 3-[3-(4'-Bromo-1,1'-biphenyl)-4-yl]-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one	0.005%
INERT INGREDIENTS	99.995%
TOTAL	100.000%

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS

HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician at once.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K intramuscularly or orally as indicated in "bromadiolone" monographs. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

Net Contents 1 1/2 OZ. (43 GM)

Mfg by



Bell Laboratories, Inc.

Madison, WI 53704 U.S.A.

EPA EST. NO. 12455 W-1

EPA REG. NO. 12455-75

DIRECTIONS FOR USE

Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other non-target animals to rodenticides. To help prevent accidents:

1. Store product out of use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in larger rodent bait stations. These stations must be resistant to destruction by signs of rat or mouse activity. Bait stations must be placed in areas where the bait must be consumed or otherwise introduced. Even though bait stations are needed in areas where the bait must be consumed, bait stations must be placed in areas where the bait must be consumed.
3. Dispose of product containers and unused bait in accordance with local, state, and federal regulations.

USE RESTRICTIONS: For use only in Norway rats, house mice and house mice.

URBAN AREAS: May be used in and around homes, residential commercial and public buildings. Contrac Place Pacs may also be used in and around structures, vehicles, ships, boats, and related structures. May not be used in fields. Do not use in sewers.

NON-URBAN AREAS: May be used in and around homes and other buildings.

Do not place bait in areas where it is possible for children, pets, domestic animals, or other non-target animals to consume the bait. Do not broadcast bait.

BEST AVAILABLE COPY

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooded livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice.

URBAN AREAS: May be used in and around the periphery of homes, industrial, commercial and public buildings. Contrac Place Pacs may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. May also be used in alleys. Do not use in sewers.

NON-URBAN AREAS: May be used inside of homes and agricultural buildings.

See right side panel for additional DIRECTIONS FOR USE.

Contrac®

RAT and MOUSE BAIT

READY TO USE PLACE PAC

KILLS RATS AND MICE

KILLS WARFARIN RESISTANT NORWAY RATS
 Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT: Bromadiolone; 3-[4-(Bromo[1,1'-biphenyl]4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one.....	0.005%
INERT INGREDIENTS	99.995%
TOTAL	100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician at once.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K, intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

Net Contents: 40 x 1 1/2 OZ.

EPA EST NO 12455-W1-1

EPA REG NO 12455-75

Mfg. by:



Bell Laboratories, Inc.
Millsboro, WI 53704 U.S.A.

DIRECTIONS FOR USE (continued from left side panel)

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS:

Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

RATS: Apply 2 to 10 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

HOUSE MICE: Place 1 Place Pac at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS - MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry area inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

ACCEPTED
MAR 30 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 12455-75