

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice infest and consume the bait. Generally, these are along with, by, or near openings, or in hidden burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Place bait in traps or areas. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

RATS: Apply 2 to 10 Place Pacs per floor room. Maintain an uninterrupted supply of fresh bait for at least 10 days. HOUSE MICE: Place 1 Place Pac at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days. MICE: Apply 1 Place Pac at 8 to 12 foot intervals. Call and dispose of all dead animals and remove bait properly. To prevent re-infestation, limit source of rodent food, water, and harborage as much as possible. If re-infestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replace as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry and inaccessible to children and pets. DISPOSAL: Do not reuse empty container. Securely wrap empty container in several layers of newspaper and discard in trash.

Contra[®]

RAT and MOUSE BAIT

READY TO USE PLACE PAC

KILLS RATS AND MICE

KILLS WARFARIN RESISTANT NORWAY RATS

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT: Bromadiolone; 3-[3-(4-Bromo-1,1-Dimethyl-4-pyridyl)-4-pyridyl]-1-phenylpropyl-4-hydroxy-2H-1-benzopyran-2-one.....	0.005%
INERT INGREDIENTS*	99.995%
TOTAL	100.000%

*Contains Dantrolen Benzoin

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep away from children, domestic animals and birds. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician at once.

NOTE TO PHYSICIAN: If applied, administer Vitamin K immediately or daily as ordered in the physician's office. Repeat as necessary based on monitoring of prothrombin level.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and aquatic life. Do not allow runoff into water.

Net Contents 1 1/2 OZ. (43 GM)

Mfg. by:



Bell Laboratories, Inc.

Moonen, PA 15704 U.S.A.

EPA EST NO 12455-W-1
EPA REG NO 12455-75

DIRECTIONS FOR USE: Is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read the entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other domestic animals to contraindications. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. Bait can be obtained from stations when trap and bait, units will be secured or otherwise immobilized. Even though bait stations are resistant to such open to baited baited, stations, bait, or other potentially destructive animals, it is each part to be replaced.
3. Dispose of product containers, and unused, spoiled, and unrecycled bait as specified on the label.

USE RESTRICTIONS: For control of Norway rats, and mice and house mice.

UNLAWFUL AREAS: May be used in and around the property of homes, industrial, commercial and public buildings. Contra Place Pacs may also be used in and around hospitals, schools, day care centers, and other public buildings.

NON-UNLAWFUL AREAS: May be used inside of homes and agricultural buildings.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not bait/label bait.

BEST AVAILABLE COPY

ACCEPTED
NOV 23 1992
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 12455-75

ACCEPTED
NOV 23 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 12455-75

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice.
URBAN AREAS: May be used in and around the periphery of homes, industrial, commercial and public buildings. Contrac Place Pacs may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. May also be used in alleys. Do not use in sewers.
NON-URBAN AREAS: May be used inside of homes and agricultural buildings.

See right side panel for additional DIRECTIONS FOR USE.

Contrac®
RAT and MOUSE BAIT

**READY TO USE PLACE PAC
 KILLS RATS AND MICE**

KILLS WARFARIN RESISTANT NORWAY RATS
 Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT: Bromadiolone; 3-[2-(4-bromo-1,1'-biphenyl)-4-yl]-3-hydroxy-1-phenylpropan-1-ylidene-2-thio-1-benzopyran-2-one 0.005%
INERT INGREDIENTS* 99.995%
TOTAL 100.000%

*Contains Dantrolene Succinate

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician at once.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K, intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin time.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

Net Contents: 40 x 1 1/2 OZ.

EPA EST. NO. 12455-WH-1
 EPA REG. NO. 12455-75

Mfg. by:



Bell Laboratories, Inc.

Madison, WI 53704 U.S.A.

DIRECTIONS FOR USE (continued from previous page)

Do not place bait in areas where there is contaminating food or surfaces that come in contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREA

Determine areas where rats or mice will consume the bait. Generally, these are gnawed openings, in or beside burrows, concealed places, between floors and where rodents or their signs have been protected from rain or snow. Remove as much as possible.

APPLICATION DIRECTIONS:

RATS: Apply 2 to 10 Place Pacs per place. Maintain an uninterrupted supply of fresh bait for at least 10 days.
HOUSE MICE: Place 1 Place Pac at each place. Maintain an uninterrupted supply of fresh bait for at least 10 days.

RATS/MICE: Replace contaminated bait as needed. Collect and dispose of all dead rodents properly. To prevent reinfestation, remove rodent food, water, and harborage as needed. If reinfestation does occur, repeat treatment. If continuous source of infestation is present, bait stations and replenish as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, inaccessible to children and pets.
DISPOSAL: Do not reuse empty container. Place original container in several layers of newspaper or cardboard in trash.