

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely feed and consume the bait. Generally, these are along walls, especially corners, in or behind furniture, in corners and under chairs, between doors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much debris as possible.

APPLICATION DIRECTIONS
RATS: Apply 2 to 10 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.
HOUSE MICE: Place 1 Place Pac at 10 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.
RATS AND MICE: Replace contaminated or soaked bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replace as needed.

STORAGE AND DISPOSAL: Store only in original container, in a dry and inaccessible to children and pets.
DISPOSAL: Do not flush empty container. Seal tightly into original container in several layers of newspaper and discard in trash.

Contrae®

RODENTICIDE

READY TO USE PLACE PAC
KILLS RATS AND MICE

KILLS WARFARIN RESISTANT NORWAY RATS
 Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT: Bromadiolone; 3-[3-(4'-Bromo-1,1'-biphenyl)-4-y]-3-hydroxy-1-phenylpropyl]; 4-hydroxy-2H-1-benzopyran-2-one.....	0.005%
INERT INGREDIENTS*	99.995%
*Contains Denatonium Benzate	TOTAL 100.000%

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician at once.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K, intramuscularly or orally as indicated in hydroquinone/acetaminophen overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds.

Mfg. by

Net Contents 1 1/2 OZ. (43 GM)



Bell Laboratories, Inc.

Madison, WI 53704 U.S.A.

EPA EST NO 12455 WI 1
 EPA REG NO 12455 76

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read the entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children, pets, domestic animals, and non-target wildlife, in a secure-resistant bait station. These stations must be resistant to tampering by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching the bait, and containers are covered and secured when they are used. Units must be secured or otherwise enclosed. Even when properly secured, bait stations are not tamper-proof.
2. Do not use bait stations in areas open to household pets, cats, dogs, birds, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, and unconsumed bait as specified on the label.

USE RESTRICTIONS: For control of Norway rats, roof rats and house mice.

URBAN AREAS: May be used in and around the company of homes, industrial, commercial and public buildings. Contrae Place Pacs may also be used in and around transport vehicles (trucks, trains, aircraft) and related parts of terminal buildings. May also be used in airports. Do not use in sewers.

NON-URBAN AREAS: May be used inside of homes and agricultural buildings.

Do not direct bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

ACCEPTED
JAN 28 1993
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered under EPA Reg. No. 12455-76

BEST AVAILABLE COPY

295

ACCEPTED
JAN 28 1993
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 12455-76

Confrac®

Rodenticide

Ready to Use Place Pac

KILLS RATS AND MICE

Kills Warfarin Resistant Norway rats

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT:

Bromadiolone; 3-[3-(4-Bromo-[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one.....0.005%
 INERT INGREDIENTS*.....99.995%

TOTAL: 100.000%

*Contains Dextrochlorpheniramine Maleate

KEEP OUT OF REACH OF CHILDREN CAUTION

(See back panel for additional precautionary statements.)

NET WT. 4 x 1 1/2 oz. PLACE PACS

EPA REG. NO. 12455-76

EPA EST. NO. 12455-WI-1

Mfg. by:



Bell Laboratories, Inc.
 Madison, WI 53704 U.S.A.

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician at once.
NOTE TO PHYSICIAN: If ingested, administer Vitamin K, intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are filled, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway rats, roof rats and house mice.

URBAN AREAS: May be used in and around the periphery of homes, industrial, commercial and public buildings. CONTRAC Place Pacs may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. May also be used in alleys. Do not use in sewers.
NON-URBAN AREAS: May be used inside of homes and agricultural buildings.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

The place pacs in this bag contain enough bait (6 oz.) to kill two to six rats.
RATS: Apply 2 to 10 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.
HOUSE MICE: Place 1 Place Pac at 8- to 12-foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.
RATS/MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry area inaccessible to children and pets.
DISPOSAL: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

BEST AVAILABLE COPY

Contract

RODENTICIDE

Ready to Use Place Pac

KILLS RATS AND MICE

KILLS WARFARIN RESISTANT NORWAY RATS

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT: Bromadiolone;
3-[3-(4'-Bromo-[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one.....0.005%
INERT INGREDIENTS*.....99.995%
TOTAL 100.000%

*Contains Denatonium Benzoate

KEEP OUT OF REACH OF CHILDREN
CAUTION

(See back panel for additional precautionary statements.)

SIX-PACK

Contains 6 — 1½ oz. Place Pacs

EPA EST. NO. 12455-WI-1

EPA REG. NO. 12455-76

Mfg. by:



Bell Laboratories, Inc.
Madison, WI 53704 U.S.A.

ACCEPTED
JAN 28 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 12455-76

BEST AVAILABLE COPY

5
495

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician at once.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K, intramuscularly or orally as indicated in biopharmaceutical monographs. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are filled, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooded livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway rats, roof rats and house mice.

URBAN AREAS: May be used in and around the periphery of homes, industrial, commercial and public buildings. CONTRAG Place Pacs may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. May also be used in areas. Do not use in sewers.

NON-URBAN AREAS: May be used inside of homes and agricultural buildings.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by ground openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

The place pacs in this bag contain enough bait (9 oz.) to kill three to nine rats.

RATS: Apply 2 to 10 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

HOUSE MICE: Place 1 Place Pac at 8- to 12-foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS/MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry area inaccessible to children and pets.

DISPOSAL: Do not reuse original container. Securely wrap original container in several layers of newspaper and discard in trash.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooled livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice.

URBAN AREAS: May be used in and around the periphery of homes, industrial, commercial and public buildings. **CONTRAC Place Pacs** may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. May also be used in alleys. Do not use in sewers.

NON-URBAN AREAS: May be used inside of homes and agricultural buildings.

See right side panel for additional DIRECTIONS FOR USE.

Contrac®

RODENTICIDE

READY TO USE PLACE PAC

KILLS RATS AND MICE

KILLS WARFARIN RESISTANT NORWAY RATS

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT: Bromadiolone; 3-β-(4-bromo-1,1'-biphenyl)-4-yl-3-hydroxy-1-phenylpropyl-4-hydroxy-2H-1-benzopyran-2-one..... 0.005%
 INERT INGREDIENTS* 99.995%
 TOTAL 100.000%

*Contains Deoxynivalenol

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician at once.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K, intramuscularly or orally as indicated in bihydrocoumatin overdose. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

Net Contents: 40 x 1/2 OZ.

EPA EST. NO. 12455-W1-1

Mfg. by:

EPA REG. NO. 12455-76



Bell Laboratories, Inc.
Mansfield, NJ 07040 U.S.A.

DIRECTIONS FOR USE (continued from left side panel)

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS:

Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

RATS: Apply 2 to 10 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

HOUSE MICE: Place 1 Place Pac at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS/MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry area inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooled livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice.

URBAN AREAS: May be used in and around the periphery of homes, industrial, commercial and public buildings. **CONTRAC Place Pacs** may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. May also be used in alleys. Do not use in sewers.

NON-URBAN AREAS: May be used inside of homes and agricultural buildings.

See right side panel for additional DIRECTIONS FOR USE.

Contrac®

RODENTICIDE

READY TO USE PLACE PAC

KILLS RATS AND MICE

KILLS WARFARIN RESISTANT NORWAY RATS

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT: Bromadiolone; 3-β-(4-bromo-1,1'-biphenyl)-4-yl-3-hydroxy-1-phenylpropyl-4-hydroxy-2H-1-benzopyran-2-one..... 0.005%
 INERT INGREDIENTS* 99.995%
 TOTAL 100.000%

*Contains Deoxynivalenol

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician at once.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K, intramuscularly or orally as indicated in bihydrocoumatin overdose. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

Net Contents: 75 x 1 1/2 OZ.

EPA EST. NO. 12455-W1-1

Mfg. by:

EPA REG. NO. 12455-76



Bell Laboratories, Inc.
Mansfield, NJ 07040 U.S.A.

DIRECTIONS FOR USE (continued from left side panel)

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS:

Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

RATS: Apply 2 to 10 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

HOUSE MICE: Place 1 Place Pac at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS/MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry area inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

ACCEPTED

JAN 28 1993
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 12455-76

BEST AVAILABLE COPY