

Reg # 12455-78

PM-14

182

ACCEPTED
NOV 9 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 12455-78

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gapped openings in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

RATS: Apply 1 to 5 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Open pac and apply 1/4 to 1/2 oz. (1-2 level teaspoons) of bait to 8- to 12-foot intervals. Larger placements (up to 2 oz) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent re-infestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL: Store only in original container, in a dry area inaccessible to children and pets. Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse empty container.

Ditrac®

Rat and Mouse Bait Place Pac

Kills Norway Rats, Roof Rats, and House Mice

ACTIVE INGREDIENT:

Diphacinone, 2-(Diphenylacetyl)-1, 3-indandione..... 0.005%


INERT INGREDIENTS..... 99.995%

TOTAL.....100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

(See additional precautionary statements on the back panel)

NET CONTENTS: 3 OZ. (85 GRAMS)

EPA EST. NO 12455-WI-1 EPA REG. NO 12455-78
 Manufactured by
 **Bell Laboratories, Inc.**
 Morristown, NJ 07954 U.S.A.

CAUTION: Keep away from humans, domestic animals, and pets. If swallowed, the material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K, intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD: Keep out of lakes, streams, or ponds.

PRECAUTIONARY STATEMENTS — HAZARD TO HUMANS AND DOMESTIC ANIMALS

READ THIS LABEL: Read the entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These bait stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are filled, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooded weasels, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway rats, roof rats and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooled livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

Place Pac Kills Norway Rats, Roof Rats, and House Mice

ACTIVE INGREDIENT:

Diphacinone; 2-(Diphenylacetyl)-1,
3-indandione..... 0.005%

INERT INGREDIENTS..... 99.995%
TOTAL..... 100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

(See additional precautionary statements on the back panel)

NET CONTENTS: 60 X 3 OZ. (85 GRAMS)

EPA EST. NO. 12455-WI-1 EPA REG. NO. 12455-78

Manufactured by:



Bell Laboratories, Inc.
MADISON, WISCONSIN U.S.A.

APPLICATION DIRECTIONS

RATS: Apply 1 to 5 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Open pac and apply 1/4 to 1/2 oz. (1-2 level tablespoons) of bait at 8- to 12-foot intervals. Larger placements (up to 2 oz) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry area inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION Keep away from humans, domestic animals, and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K₁ intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD: Keep out of lakes, streams or ponds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooled livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

Ditrac® Rat and Mouse Bait

Place Pac Kills Norway Rats, Roof Rats, and House Mice

ACTIVE INGREDIENT:

Diphacinone; 2-(Diphenylacetyl)-1,
3-indandione..... 0.005%

INERT INGREDIENTS..... 99.995%
TOTAL..... 100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

(See additional precautionary statements on the back panel)

NET CONTENTS: 20 X 3 OZ. (85 GRAMS)

EPA EST. NO. 12455-WI-1 EPA REG. NO. 12455-78

Manufactured by:



Bell Laboratories, Inc.
MADISON, WISCONSIN U.S.A.

SELECTION OF TREATMENT AREAS:

Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along wall, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

RATS: Apply 1 to 5 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Open pac and apply 1/4 to 1/2 oz. (1-2 level tablespoons) of bait at 8- to 12-foot intervals. Larger placements (up to 2 oz) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry area inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

292