

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE
	12455-84	MAR 23 1993
	TERM OF ISSUANCE	
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREГИSTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	NAME OF PESTICIDE PRODUCT	
	DITRAC RODENTICIDE READY-TO-USE PLACE PAC	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Bell Laboratories, Inc.
 3699 Kinsman Boulevard
 Madison, WI 53704

5436748 $\frac{165}{17}$

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Add the phrase, "EPA Registration No. 12455-84" to your label before you release the product for shipment.
3. Submit one copy of final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

This registration will be subject to cancellation in accordance with FIFRA sec. 6(e) if you do not comply with these conditions. Your release for shipment of the product constitutes acceptance of these conditions.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL: *[Signature]* DATE: _____

A stamped copy of the label is enclosed for your records.

RAF

Robert A. Forrest
Product Manager (14)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosures: 1) Stamped label
2) A-79 Enclosure

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely feed and consume bait. Generally, these are dry areas, by ground openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Place bait from rats or mice. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

RAIS: Apply 2 to 10 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

HOUSE MICE: Place 1 Place Pac at 1 to 1 1/2 feet intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RAIS AND MICE: Replace contaminated or soaked bait immediately. Collect and dispose of all dead animals and remove bait promptly. To prevent reinfestation, have sources of rodent, bait, water, and food kept as much as possible. If reinfestation does occur, repeat treatment. There is continuous action of Diphacinone to prevent, establish permanent bait stations and replace as needed.

STORAGE AND DISPOSAL:

STORAGE: Store only in original container, in a dry area inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Secondary or original container if level of liquid is near paper and discard in 7 days.

Ditraac®

Rodenticide
READY TO USE PLACE PAC

**Kills Norway Rats, Roof Rats,
and House Mice**

ACTIVE INGREDIENT: Diphacinone;

2-(Diphenoxyacetyl)-1,3-indandione	0.005%
INERT INGREDIENTS	99.995%
TOTAL	100.000%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUTIONARY STATEMENTS


HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CANTON: Keep away from humans, domestic animals and pets. If swallowed, the material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K intramuscularly or orally, as indicated in the hydroncourse-overdose. Repeat as necessary based on monitoring of prothrombin time.

ENVIRONMENTAL HAZARD: Keep out of lakes, streams or ponds.

Net Contents 1 1/2 OZ. (43 GM)

Mfg. by:  Bell Laboratories, Inc.
Madison, WI 53704 U.S.A.

EPA EST NO 12455-W-1
EPA REG NO 12455-04

DIRECTIONS FOR USE This is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- READ THIS LABEL:** Read the entire label and follow all use directions and use precautions.
- IMPORTANT:** Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:
1. Store product out of reach of children and pets.
 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in larger areas, or in larger areas, or in larger areas, or in larger areas. These rodents must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching the bait compartments and obtaining bait. Bait can be eaten from children when they are used with mice. Bait must be secured in appropriate household. Each storage bait station are located in areas open to household livestock, raccoons, bears, or other primary ungulate animals, or in areas prone to vandalism.
 3. Observe all product cautions, and unusual, spilled, and unreturned bait as specified on this label.

USE RESTRICTIONS:
For control of Norway rats, roof rats and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or food areas that come in direct contact with food. Do not broadcast bait.

ACCEPTED
MAR 23 1993
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 12455-84

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Keep away from humans, domestic animals, and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K₁ intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD: Keep out of lakes, streams or ponds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

Ditrac® Rodenticide

READY TO USE PLACE PAC


KILLS RATS AND MICE

ACTIVE INGREDIENT: Diphacinone:	
2-(Diphenylacetyl)-1,3-indandione	0.005%
INERT INGREDIENTS	99.995%
TOTAL	100.000%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Net Contents: 120 x 1 1/2 OZ. (43 GM)

Mfg. by:

 **Bell Laboratories, Inc.**
Madison, WI 53704 U.S.A.

EPA REG. NO. 12455-84 EPA EST. NO. 12455-WI-1

ACCEPTED
MAR 23 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 12455-84

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS:

Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

RATS: Apply 2 to 10 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.
HOUSE MICE: Place 1 Place Pac at 8- to 12-foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.
RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry area inaccessible to children and pets.
DISPOSAL: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

BEST AVAILABLE COPY

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K₁ intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD: Keep out of lakes, streams or ponds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

Ditrac® Rodenticide

READY TO USE PLACE PAC


KILLS RATS AND MICE

ACTIVE INGREDIENT: Diphacinone:	
2-(Diphenylacetyl)-1,3-indandione	0.005%
INERT INGREDIENTS	99.995%
TOTAL	100.000%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Net Contents: 35 x 1 1/2 OZ. (43 GM)

Mfg. by:

 **Bell Laboratories, Inc.**
Madison, WI 53704 U.S.A.

EPA REG NO 12455-84 EPA EST NO 12455-WI-1

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS:

Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

RATS: Apply 2 to 10 Place Pacs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.
HOUSE MICE: Place 1 Place Pac at 8- to 12-foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.
RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry area inaccessible to children and pets.
DISPOSAL: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.