

R. P. KNISELY CO.

17126 Detroit Ave.

Cleveland, Ohio 44107

NET WT. _____ LBS.

MICROBIOCIDE-K

APPLICATION

Microbiocide-K is used to control algae, bacteria and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime and other deposits. An initial slug addition of 3.9 to 9.75 fluid ounces of Microbiocide-K per 1000 gallons of water, to provide a concentration of 32 to 80 parts per million of Microbiocide-K based on the total weight of water in the system is recommended. Repeat until control is achieved.

Subsequent slug additions of 1.0 to 9.75 fluid ounces of Microbiocide-K per 1000 gallons of water (8 to 80 parts per million of Microbiocide-K should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of the cooling tower.

COMPOSITION

	Percent by weight
<i>Active ingredient</i>	15.0%
<i>Poly (oxyethylene (dimethyliminio) ethylene- (dimethyliminio) ethylene dichloride)</i>	15.0%
<i>Inert ingredient</i>	85.0%
<i>Weight per gallon</i>	8.58 pounds
<i>Weight of active ingredient per gallon</i>	1.29 pounds

EPA Reg. No. 12459-1

CA

KEEP OUT OF THE HANDS OF CHILDREN.
Harmful if swallowed and eyes. In case of contact with skin or eyes, wash with plenty of water for at least 15 minutes. If irritation persists get medical attention. Avoid contamination of food.

This product is toxic to fish. Do not use in or near streams, rivers, ponds, lakes, or other bodies of water. Do not contaminate water, food, or feed. Do not use on lawns, lawns, or other areas where people or animals may be present. Do not contaminate equipment or disposal containers. See directions directed on label. Destroy when empty.

ACCEPTED

March 2, 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION

NET WT. _____ LBS.

MICROBIOCID-K

N

algae, bacteria
al and industrial
e, systems must
microbiological
al slug addition
icrobiocide-K per
a concentration
Microbiocide-K
r in the system
trol is achieved.

to 9.75 fluid
gallons of water
biocide-K should
as needed. The
on the relative
ty of the micro-
should be made

COMPOSITION

	Percent by weight
<i>Active ingredient</i>	15.0%
<i>Poly (oxyethylene (dimethyliminio) ethylene-</i> <i>(dimethyliminio) ethylene dichloride)</i>	15.0%
<i>Inert ingredient</i>	85.0%
<i>Weight per gallon</i>	8.58 pounds
<i>Weight of active ingredient per gallon</i>	1.29 pounds

EPA Reg. No. 12459-1

CAUTION

KEEP OUT OF THE REACH OF CHILDREN
Harmful if swallowed. Avoid contact with skin and eyes. In case of contact flush with plenty of water for at least 15 minutes. If eye irritation persists get medical attention. Avoid contamination of food.

This product is toxic to fish. Do not add treated effluent to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste. Use only as directed on label. Do not reuse container. Destroy when empty.

ACCEPTED

March 2, 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION