



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Antimicrobials Division (7510P)  
 1200 Pennsylvania Ave., N.W.  
 Washington, D.C. 20460

EPA Reg. Number:

12487-20

Date of Issuance:

3/24/2021

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
 (under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

MicroControl 2525

Name and Address of Registrant (include ZIP Code):

National Colloid Company  
 906 Adams Street  
 Steubenville, OH 43952

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Antimicrobials Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

John Hebert, Branch Chief  
 Regulatory Management Branch I,  
 Antimicrobials Division (7510P)  
 Office of Pesticide Programs

Date:

March 24, 2021

2. You are required to comply with the data requirements described in the DCI Order identified below:
  - a. GDCI- 0989951-1626      EPA-HQ-OPP- 2015-0736
  - b. GDCI -107103-1709      EPA-HQ-OPP- 2013-0605
  - c. GDCI -107104-1479      EPA-HQ-OPP- 2013-0605

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Reevaluation Team Leader (Team 36): <http://www2.epa.gov/pesticide-contacts/contacts-office-pesticide-programs-antimicrobial-division>

3. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 12487-20.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated March 24, 2021

If you have any questions, please contact Zebora Johnson by phone at (703) 308-7080, or via email at [johnson.zeboraj@epa.gov](mailto:johnson.zeboraj@epa.gov).

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER.** Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed, absorbed through the skin or inhaled. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist. Wear goggles or face shield, chemical-resistant gloves, long-sleeve shirt, long pants, shoes, socks and chemical-resistant apron when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

### ENVIRONMENTAL HAZARD

This product is toxic to fish, aquatic invertebrates, oysters and shrimp. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### WATER TREATMENT

(Not for use in CA)

Do not use water containing residues from use of this product to irrigate crops for food or feed.

**AIR WASHERS, INDUSTRIAL AND/OR COMMERCIAL RECIRCULATING COOLING WATER TOWERS, RETORT WATER SYSTEMS, EVAPORATIVE CONDENSERS, HEAT EXCHANGE TRANSFER WATER SYSTEMS, DAIRY SWEETWATER SYSTEMS, HYDROSTATIC STERILIZERS, PASTEURIZERS AND WARMERS:** For best results, clean heavily contaminated systems before treatment with this product. If soap or anionic detergent is used, rinse thoroughly before charging with this algicide. {Cooling tower waters that are inherently low in algae growth and bacteria count may be adequately controlled by the lower range of these dosages.} Repeat every seven days or increase frequency if needed. Should slime develop again, repeat initial dosage. This product is used only in industrial air washers and air washer systems which have mist-eliminating components.

- Dosing Location:** This product is to be applied at a point in the system where it will be uniformly mixed, such as the basin area, the sump, or another reservoir or collecting area.

SEE ADDITIONAL SHEET FOR OTHER DIRECTIONS FOR USE.

# MicroControl 2525

{All text in brackets [xxx] is optional and may or may not be intended on a final label.}

### ACTIVE INGREDIENTS

Glutaraldehyde ..... 25.7%  
Didecyl dimethyl ammonium chloride ..... 7.5%  
Alkyl (50% C<sub>14</sub>, 40% C<sub>12</sub>, 10% C<sub>16</sub>)  
dimethyl benzyl ammonium chloride..... 5.0%

**OTHER INGREDIENTS** ..... 61.8%  
**TOTAL** ..... 100.00%

Weight Approx. 8.56 lbs./gal.

## DANGER

### KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

**[Note to Reviewer:** In accordance with 40 CFR 156.68(d), all first aid statements, as prescribed, will appear on the front panel of the product label.]

### FIRST AID

In case of emergency, call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

**IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

**IF SWALLOWED:** Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Call a poison control center or doctor immediately for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. For chemical and medical or environmental emergencies, call INFOTRAC at (800) 535-5053 24 hours a day, 7 days a week.



The National Colloid Company  
906 Adams St.  
Steubenville, OH 43952  
(740) 282-1171

EPA Reg. No. 12487-E  
EPA Est. No. 12487-OH-01

Net Contents: \_\_\_\_\_  
Lot No.: \_\_\_\_\_

Made in the USA

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

### PESTICIDE STORAGE

Store only in original container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

### PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING

Non-Refillable Container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Fill the container 1/4 full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

### WARRANTY STATEMENT

Read product Safety Data Sheet prior to use. PRODUCT WARRANTY, DISCLAIMER, AND LIMITATION OF LIABILITY ARE FOUND on the product Safety Data Sheet. Unless inconsistent with applicable law, use of Product signifies agreement with these provisions

**ACCEPTED**

03/24/2021

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 12487-20

2. **Dosing Conditions:** This product must be applied when the system is in jeopardy of being affected or after cleaning systems where efficiency is already impaired. Tower bleed off valves must be closed to permit a retention time of 4 hours.

3. **Method of Application:**

a. **INTERMITTENT OR SLUG METHOD**

**Initial Dose:** When the system is noticeably fouled, apply 17 – 34 oz. of this product per 1,000 gal. of water (50 – 100 ppm active) in the system. Repeat until control is achieved.

**Subsequent Dose:** When microbial control is evident, add 7 – 17 oz. of this product per 1,000 gal. of water (20 – 50 ppm active) in the system weekly or as needed to maintain control.

b. **CONTINUOUS FEED METHOD**

**Initial Dose:** When the system is noticeably fouled, apply 17 – 34 oz. of this product per 1,000 gal. of water (50 – 100 ppm active) in the system.

**Subsequent Dose:** Maintain this treatment by starting a continuous feed of 4 – 17 oz. of this product per 1,000 gal. of water (10 – 50 ppm active) lost by blowdown.

**AUXILIARY SYSTEMS AND SERVICE WATER:** Add 17 – 34 oz. of this product per 1,000 gal. of water (50 – 100 ppm active) in the system continuously. This product must be added to the system at a point of uniform mixing by slug or intermittent feed or by spraying onto a waste pile. The frequency of feed or spray and the duration of treatment will depend upon the severity of the contamination. Additions to water systems must be made during the pumping operation and as close to the pump as possible to ensure adequate mixing.

**INDUSTRIAL WASTEWATER SYSTEMS: Wastewater Systems, Wastewater sludge, and Wastewater Holding Tanks:** This product is added to a wastewater system or sludge at a convenient point of uniform mixing such as digester. Add 85 – 420 oz. of this product per 1,000 gal. of wastewater or sludge (250 – 1250 ppm active).

**OIL FIELD, GAS PRODUCTION, AND TRANSMISSION PIPELINE SYSTEMS**

(Not for use in CA)

**OIL FIELD, GAS PRODUCTION, TRANSMISSION PIPELINE, AND SYSTEMS:** Specific treatment requirements vary among oil and/or gas field sites and subsystem components. Oil field fluids and subsystems most commonly requiring microbial contamination control are raw water sources, separators, ballasts, storage and mixing tanks, screens, surface injection equipment, production equipment (such as injection and production piping casing, completion, and valving) and the formation itself. The primary point of treatment will vary among oil and/or gas field operations depending on the site problems, water-flood treatment methods and equipment. This product must be added where it will disperse rapidly and uniformly to the desired area of treatment.

Additions of this product must be made with the proper type of metering pump equipment, suction (low pressure) side of pumping equipment or similar device. This product must be added to the system by slug, continuous, or on an intermittent basis, depending on the degree of system fouling.

**OIL FIELD WATER FLOOD SYSTEMS AND FRACTURING FLUIDS:** This product must be added to the water flood water disposal system at a point of uniform mixing.

1. **Continuous Use:** Add 1.3 – 65 gal. of this product per 10,000 gal. of flood water (50 – 2500 ppm active) to control slime forming and sulfate reducing bacteria. Levels for effective control will vary depending on conditions at the site.

2. **Intermittent Use:** Add at a rate of 1.3 – 65 gal. of this product per 10,000 gal. of flood water (50 – 2500 ppm active) for 4 – 8 hours per day, one to four times a week as needed to maintain control.

3. **Treatment of flow back return water (Post Hydraulic Fracturing):** Dose at a rate of 1.3 – 65 gal. of this product per 10,000 gal. of water of flood water {(50 – 2500 ppm active)} for 4 – 8 hrs. per day, one to four times a week as needed to maintain control.

**FRACTURING FRAC FLUIDS:** Add this product to the frac water storage tanks or directly into the well head injection pipeline as the water is being pumped down-hole.

**Dose Range:** Add 1.3 – 65 gal. of this product per 10,000 gal. of flood water (50 – 2500 ppm active) to control slime forming and sulfate reducing bacteria. Levels for effective control will vary depending on conditions at the site.

**OIL AND GAS PRODUCTION AND TRANSMISSION PIPELINES AND SYSTEMS:** For the control of sulfate-reducing bacteria and slime forming bacteria, this product must be added to a gas production or transmission pipeline via direct injection at a point where uniform and maximum distribution will occur. The application must be conducted to ensure maximum distribution of the product through the internal surface of the pipeline by adding an amount of biocide which eventually comes out the other end of the pipeline. Criteria for success of the treatment will be reduction in bacterial count and/or corrosion rates. To facilitate application, it is desirable to dilute the product with an appropriate solvent immediately before use. The concentration in the solvent must not fall below an active concentration range of 500 – 5000 ppm active based on the volume of water in the pipeline. Injections to the system must be weekly, or as needed to maintain control.

**GAS STORAGE WELLS AND SYSTEMS:** Treat individual injection wells with 0.6 – 6.5 gal. of this product per 1,000 gal. of water (250 – 2500 ppm active). Update treatment rate as needed. This product must be diluted by the water present in the formation. Injection takes place before gas is injected and may be repeated yearly or as needed to maintain control.

Individual drips should be treated with a sufficient quantity of this product to produce a concentration of 100-1000 ppm active of this product when diluted by the water present in the drip. Injections should be repeated yearly or as needed to maintain control.

**PIPELINE PIGGING AND SCRAPING OPERATIONS:** Add this product to slug water immediately following the scraper (keep the water volume to a minimum and contained between the scraper and the following trailing pig). Add an effective concentration of .13 – 1.31 gal. of product per 1,000 gal. of water (500 – 5000 ppm active) depending on the length of the pipeline and the severity of the biofouling.

**DRILLING, COMPLETION AND WORKOVER FLUIDS SYSTEMS:** This product is to be added to these fluid systems at a point of uniform mixing, such as a circulating, holding or mud tank. Levels for effective control will vary depending on conditions at the site and the severity of the contamination.

1. **Initial treatment:** Add 0.6 – 13 gal. of this product per 1,000 gal. of freshly prepared fluid (25 – 500 ppm active).
2. **Maintenance dosage:** Add 0.6 – 13 gal. of this product per 1,000 gal. of freshly prepared fluid (25 – 500 ppm active).

**PACKER FLUIDS:** This product is to be added to the packer fluid at a point of uniform mixing such as a circulating holding tank and other mixing device locations. Add 0.6 – 8 gal. of this product per 10,000 gal. of freshly prepared packer fluid (25 – 300 ppm active). Levels for effective control vary depending on conditions at the site and the severity of contamination. Seal the treated packer fluid in the wall between the casing and the production tube.

**HYDROTESTING:** Treat water used to hydrotest pipelines or vessels by adding 0.3 – 5 gal. of this product per 1,000 gal. of water (50 – 2000 ppm active) depending on the water quality and length of time the equipment will remain idle.

**PULP AND PAPER MILLS**

(Not for use in CA)

Do not use to treat paper or paperboard which will contact food.

**SLIMICIDE APPLICATIONS:** Do not use to treat paper or paperboard which will contact food. Apply this product to the paper making system at a point of uniform mixing such as, thin or thick stock chests, save-all tank, process tank or whitewater tank.

**Initial Dose:** When system is noticeably contaminated, add 0.5 – 53 gal. of this product per 100,000 gal. of whitewater (1 – 200 ppm active) to be treated as a continuous or slug dose. Repeat until control is achieved. Heavily fouled systems must be boiled out prior to initial treatment.

**Subsequent Dose:** When microbial control is evident, add 0.5 – 53 gal. of this product per 100,000 gal. of whitewater (1 – 200 ppm active) to be treated as necessary to maintain control.

**WATER-BASED COATING, PIGMENTS AND FILLER SLURRIES FOR PULP AND PAPERBOARD:** Application of this product must be made at a point in the system where mixing action is good or can be made at the size press or water box.

**Dosing Application:** Apply at a rate of 0.13 – 0.79 lbs. per 1,000 lbs. of dry powder or 0.065 – 0.393 kg of this product per metric ton of dry slurry to produce a concentration of 130.9 – 785.3 ppm active as product (based on slurry solids) in the mixed slurry.